

Delayed hypoglycemic effect of insulin overdose in a diabetic child: a case report

Khalid Al Noaim^{1,2*}, Amir Babiker^{2,3,4}, Angham Al Mutair^{2,3,4},
Mohsen Al Atawi^{2,3,4}, Mohammed Al Dubayee^{2,3,4},
Fahad Al Juraibah^{2,4}, Ibrahim Al Alwan^{3,4}

European Journal of
Medical Case Reports

Volume 3(2):91–94

© EJMCr. <https://www.ejmcr.com/>

Reprints and permissions:

<https://www.discoverpublish.com/>

<https://doi.org/10.24911/ejmcr/>

173-1556259277



ABSTRACT

Background: Insulin overdose can result in fatal hypoglycemia. We report on a diabetic child who received an unintentional insulin overdose.

Case Presentation: A 13-year-old girl with type 1 diabetes, who was on insulin pump therapy, developed high blood glucose (BG) level due to pump malfunction. She gave herself 7 ml (700 units) instead of seven units by an error. She presented to hospital 2 hours later when this was recognized. The initial BG in the emergency room was 13.9 mmol/l. A dextrose infusion was started 6 hours after the insulin injections when her BG dropped significantly before it reached a lowest level of 3.3 mmol/l, 9 hours after the event. It was only possible to discontinue this infusion 15 hours after insulin injections when BG was stable.

Conclusion: Hypoglycemia could be delayed in insulin overdose in diabetic children. This may necessitate prolonged monitoring beyond the usual half-life of insulin.

Keywords: Insulin, overdose, delayed hypoglycemia, dextrose, glucagon, aspart, diabetes.

Received: 29 April 2018

Accepted: 07 May 2019

Correspondence to: Khalid Ibrahim Al Noaim

*College of Medicine, King Faisal University, Al Ahsa, Saudi Arabia.

Type of Article: CASE REPORT

Specialty: Pediatric Endocrinology

Email: kalnoaim@kfu.edu.sa; n3aaim@hotmail.com

Funding: None

Full list of author information is available at the end of the article.

Declaration of conflicting interests: None

Background

Insulin overdose can be a significant emergency condition due to risk of fatal hypoglycemia [1]. Mortality accounts for 2% of the cases of insulin overdoses [2]. There are few reports that discuss unintentional insulin overdose in teenagers as most of the reported cases were secondary to mental disorders or suicidal attempts [2,3]. Insulin aspart is rapid acting insulin with potential of short duration hypoglycemic episodes [4]. However, the hypoglycemic effect of insulin can be prolonged in case of insulin overdose with maximum effect at the first few hours after the injection(s) [5]. Different hypotheses were postulated to explain the pathophysiological process of this prolonged action of insulin; some were confirmed by the experiments [6]. We report on a teenager with type 1 diabetes mellitus (T1DM) who received an unintentional insulin aspart overdose, which resulted in a delayed onset of hypoglycemia.

Case Presentation

We report a 13-year-old Saudi girl, who is a known case of T1DM on insulin pump therapy. She developed blood glucose (BG) reading “HIGH” which is more than 33.3 mmol/l (600 mg/dl) due to pump malfunction. Mistakenly, instead of taking seven units of insulin aspart as a correction dose, she gave herself 7 ml (700 units) of insulin aspart by insulin syringes as been advised on the dose over the phone by her sister, who is a doctor,

and usually counsel her younger sister in management. She received seven injections (1 ml/syringe) that were given in the patient abdomen and both thighs. The patient presented to our Emergency Department after 2 hours from the injection time when this error was recognized. She was fully conscious with normal vital signs. The injection site in her abdomen was lipodystrophic. The initial BG reading at the emergency room was 13.9 mmol/l (250 mg/dl), and that falsely reassured the emergency room (EM) staff. Therefore, the patient was not started on intravenous (IV) dextrose infusion, and her insulin pump was suspended. However, the patient continues on frequent BG monitoring every 30 minutes. Her oral intake was minimal due to poor appetite. Her BG was decreased gradually till it reached 3.8 mmol/l (70 mg/dl) after 6 hours from the received injections, hence, started on 10% dextrose IV infusion. After that, BG improved but started to decline again gradually during the dextrose infusion reaching the lowest level of 3.3 mmol/l (60 mg/dl), after 9 hours of the insulin injections. This was treated by bolus of 10% dextrose and 1 mg of intramuscular glucagon (Figure 1). Investigations, ordered later in the patient by endocrinology team, showed equivalent levels of BG in the lab, normal electrolytes, venous blood gas, and renal functions (Tables 1 and 2). Six hours later, i.e., 15 hours after the overdose by insulin injections, dextrose infusion was discontinued due to stabilization of BG




Figure 1. Hospital BG levels over time from insulin injections.

Table 1. Initial level of electrolytes at presentation.

ELECTROLYTES	LEVEL
Na (mmol/l)	132
K (mmol/l)	3.6
Cl (mmol/l)	103
BUN (mmol/l)	5
Creatinine (µmol/l)	79

Table 2. Level of C-peptide and insulin (at 12 hours post-injection).

Insulin (uIU/ml)	3.8
C-Peptide (ng/ml)	0.5

level. Nevertheless, the patient was kept on continuous monitoring till completed 12 hours, in which she did not develop hypoglycemia, after discontinuation of dextrose infusion.

Discussion

Insulin overdose can be a serious condition [1]. In a recent report, T1DM patients accounted for more than 50% of insulin overdose cases [2]. An unintentional insulin overdose accounts for 5% of the total cases. Approximately, 50% of the total cases had psychiatric illnesses [2]. The recovery from risk of hypoglycemia following insulin overdose occurs maximally in 1 week with a mean time of approximately 72 hours [2,5].

It is important to identify the maximum glucose clearance rate in cases of insulin overdose as this could anticipate the development of hypoglycemia and the duration needed for glucose infusion [7–9]. In our case, the initial BG reading at home in the glucometer was “HIGH” which means more than 33.3 mmol/l (600 mg/dl), but the exact high level is unknown. Therefore, the rate of decreasing of glucose during the initial 2 hours after the injections could not be accurately identified. Although some reported cases had a delayed onset of hypoglycemia, after a huge dose of insulin injection [9,10].

Teenager’s behavior and problems related to insulin injection site are well known factors of diabetic mismanagement [11]. The other possibility that could explain the

situation in our patient is that our patient might have not actually received the claimed insulin aspart dose due to either expired insulin, poor technique of injection, or a part of teenager’s manipulation. Last, but not the least, the patient could have used lipohypertrophy site of injection that consequently leads to delayed of insulin absorption.

The prolonged effect of hypoglycemia in insulin overdose is more commonly dose rather than insulin type dependent [5]. Different mechanisms might explain the prolonged hypoglycemic effect in short acting insulin overdose. First, decrease absorption of insulin from the injection site due to the mechanical effect that leads to compression in the local microvascular circulation, which is supported by experimental studies [5,12]. In addition, surgical incisions of the injection site have been used as a modality of treatment [13]. Second, prolongation of half-life of insulin due to decreased insulin clearance when the maximum capacity of hepatic and renal receptors of insulin excretion is exceeded [6]. Moreover, insulin pharmacokinetics is suggested to be unique. For instance, the IV administration of glargine, long acting insulin, did not result in hypoglycemia in compared to similar dose given by subcutaneous route [14].

The severity and serious consequences are more common in non-diabetic patient who received the insulin overdose, because of the possible presence of insulin autoantibodies as well as absence of endogenous insulin in patient with T1DM [15]. Nonetheless, the most common electrolyte imbalance, other than hypoglycemia, is hypokalemia due to insulin-induced potassium shift from extracellular to intracellular compartment [2].

There are different modalities of treatments of insulin overdose, and the most common used is an IV glucose therapy, which includes boluses and infusions. Other modalities include glucagon, oral carbohydrates, surgical incision at the injection site, octreotide, and hydrocortisone [2,9,13]. In addition, other management should be include electrocardiogram (ECG) monitoring and optimization of serum potassium level that could be drop secondary to insulin overdose. Use of continues glucose monitoring device during the hospitalization may be helpful to guide the physician when to stop the treatment [16].

Understanding the possibilities of progress in different scenarios, as above, in term of immediate or slow development of hypoglycemia is a cornerstone in management of such cases. In our patient, the impression of stability of BG initially lead to the false opinion that possibly delayed the initiation of IV glucose infusion and the requesting insulin level and c-peptide in the list of investigations at presentation. Using of insulin pen instead of insulin syringe may decrease the risk of miscalculation of insulin correction dose.

Conclusion

Insulin overdose may lead to a delayed hypoglycemic effect. Hence, affected patients should be monitored for this risk even beyond the half-life of the overdosed insulin.

Acknowledgment

The author would like to thank the patient and her family for agreeing this case report to be published.

List of Abbreviations

BG	Blood glucose
ECG	Electrocardiogram
ER	Emergency room
IV	Intravenous
T1DM	Type 1 diabetes mellitus

Consent for publication

Informed consent was obtained from the patient and her family.

Ethical approval

The manuscript follows the rules and regulation of institution review board in the King Abdullah International Medical Research Center (KAIMRC) in publishing of patient information including confidentiality and consent for publication.

Author details

Khalid Al Noaim^{1,2}, Amir Babiker^{2,3,4}, Angham Al Mutair^{2,3,4}, Mohsen Al Atawi^{2,3,4}, Mohammed Al Dubayee^{2,3,4}, Fahad Al Juraibah^{2,4}, Ibrahim Al Alwan^{3,4}

1. College of Medicine, King Faisal University, Al Ahsa, Saudi Arabia
2. Department of Pediatrics, Ministry of the National Guard—Health Affairs, Riyadh, Saudi Arabia
3. College of Medicine, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia
4. King Abdullah International Medical Research Center, Riyadh, Saudi Arabia

References

1. Batalis N, Prahlow J. Accidental insulin overdose. *J Forensic Sci.* 2004;49(5):1–4. <https://doi.org/10.1520/jfs2004059>
2. Johansen N, Christensen M. A systematic review on insulin overdose cases: clinical course, complications and treatment options. *Basic Clin Pharmacol Toxicol.* 2018;122(6):650–9. <https://doi.org/10.1111/bcpt.12957>
3. Arem R, Zoghbi W. Insulin overdose in eight patients. *Medicine.* 1985;64(5):323–32. <https://doi.org/10.1097/00005792-198509000-00004>
4. Adams G, Meal A, Morgan P, Alzahrani Q, Zobel H, Lithgo R, et al. Characterisation of insulin analogues therapeutically available to patients. *PLoS One.* 2018;13(3):e0195010. <https://doi.org/10.1371/journal.pone.0195010>
5. Ohyama T, Saisho Y, Muraki A, Kawai T, Itoh H. Prediction of recovery time from hypoglycemia in patients with insulin overdose. *Endocr J.* 2011;58(7):607–11. <https://doi.org/10.1507/endocrj.k11e-018>
6. Sato Y, Mizuno Y, Suganuma K, Shioto K, Ikeda T, Yamashita K, et al. Pharmacokinetics of insulin disappearance after massive overdosing. *Endocr J.* 2018;65(11):1147–53. <https://doi.org/10.1507/endocrj.ej18-0118>
7. Fasching P, Roden M, Stühlinger H, Kurzemann S, Zeiner A, Waldhäusl W, et al. Estimated glucose requirement following massive insulin overdose in a patient with type 1 diabetes. *Diabetic Med.* 1994;11(3):323–5. <https://doi.org/10.1111/j.1464-5491.1994.tb00279.x>
8. Bizzotto R, Natali A, Gastaldelli A, Muscelli E, Krssak M, Brehm A, et al. Glucose uptake saturation explains glucose kinetics profiles measured by different tests. *Am J Physiol Endocrinol Metab.* 2016;311(2):E346–57. <https://doi.org/10.1152/ajpendo.00045.2016>
9. Tariq K, Tariq S, Denney Queen A. Role of steroids in refractory hypoglycemia due to an overdose of 10,000 units of insulin glargine: a case report and literature review. *AACE Clin Case Rep.* 2018;4(1):e70–4. <https://doi.org/10.4158/ep171780.cr>
10. Mork T, Killeen C, Patel N, Dohnal J, Karydes H, Leikin J. Massive insulin overdose managed by monitoring daily insulin levels. *Am J Therap.* 2011;18(5):e162–6. <https://doi.org/10.1097/mjt.0b013e3181f4eadb>
11. Borus J, Laffel L. Adherence challenges in the management of type 1 diabetes in adolescents: prevention and intervention. *Curr Opin Pediatr.* 2010;22(4):405–11. <https://doi.org/10.1097/mop.0b013e32833a46a7>
12. Hildebrandt P, Birch K, Sestoft L, Vqlund A. Dose-dependent subcutaneous absorption of porcine, bovine and human NPH insulins. *Acta Med Scand.* 2009;215(1):69–73. <https://doi.org/10.1111/j.0954-6820.1984.tb04971.x>
13. Droste J, Hundia V, Pettit A, Narayan N, Nejm A. Excision of injection site substantially reduced serum insulin concentration in a potentially life-threatening insulin analogue overdose. *Prac Diabetes.* 2012;29(6):243–5. <https://doi.org/10.1002/pdi.1700>
14. Thornton S, Gutovitz S. Intravenous overdose of insulin glargine without prolonged hypoglycemic effects. *J Emerg Med.* 2012;43(3):435–7. <https://doi.org/10.1016/j.jemermed.2011.06.038>
15. Stapczynski JS, Haskell RJ. Duration of hypoglycemia and need for intravenous glucose following intentional overdoses of insulin. *Ann Emerg Med.* 1984;13(7):505–11. [https://doi.org/10.1016/s0196-0644\(84\)80513-2](https://doi.org/10.1016/s0196-0644(84)80513-2)
16. El-Laboudi AH, Misra S, Martineau M, Deol P, Sanders A, Oliver N. Intentional large insulin overdose captured on a continuous glucose monitor. *J Diabetes Sci Technol.* 2015;9(4):929–31. <https://doi.org/10.1177/1932296815579691>

Summary of the case

Patient (gender, age)	1	Female, 12 years
Final diagnosis	2	T1DM, insulin overdose
Symptoms	3	Hypoglycemia
Medications	4	IV 10% dextrose, IM glucagon
Clinical procedure	5	N/A
Specialty	6	Endocrinology