







**Figure 1.** 3-year-old girl with normal physique who suffered blunt abdominal trauma and was diagnosed to be having pancreatic pseudocyst during follow-up.




**Figure 2.** Plain thoraco-abdominal skiagram shows normally placed structures with normal bowel gas shadows pattern. No air-fluid level seen, to suggest any intestinal obstruction.




**Figure 3.** Ultrasound images. (a) Axial section shows fluid collection (white star) anterior to pancreas and SP. This also showed some echogenic debris (green arrow). (b) Oblique scan shows total extent of the collection region (white star) with the mobile echogenic sediments (green arrow). SP spine.




**Figure 4.** Color flow imaging. (a) Anechoic collection (white star) anterior to the superior mesenteric artery (green arrow) and pancreatic body region being on posterior side (white arrow). (b) Oblique view shows displayed spleno-portal axis (white arrow) by the anechoic fluid collection (white star).




**Figure 5.** Contrast enhanced computerized tomography abdomen. (a) Axial section shows well-defined collection (white arrow) adjacent to stomach (s). (b) Sagittal section shows posterior placement of pseudocyst (red star). (c) Coronal section shows pseudocyst (red star) placed in between the liver and stomach (S).



**Figure 6.** Magnetic resonance imaging images (a) T1W axial section shows hypointense collection (inverted arrow) with adjoining stomach (red star). (b) T2W axial section shows the same collection as hyperintense (black star) in-between GB and ST. SPL lies posterolateral to the stomach. (c) Coronal T2W image shows well-delineated hyperintense pseudocyst postero-medial to the ST. GB, gall bladder; ST, stomach; SPL, Spleen.



**Figure 7.** Magnetic resonance imaging images contd. (a) Axial section of BTFE sequence shows hyperintense pseudocyst (green star) with head of pancreas (white arrow) and spleno-portal axis (red arrow). (b) Coronal BTFE shows pseudocyst (red star) with the stomach on the left-hand side (green star) and the gall bladder on the right-hand side (green arrow). BTFE, Balanced turbo-field-echo.



**Figure 8.** Magnetic resonance cholangio-pancreaticography image. There is a common normal entry of the common bile and pancreatic ducts (white arrow). Hyperintense well-defined pseudocyst (red star) lies in between the gall bladder (G) and the stomach (S). The child was diagnosed as having pseudocyst formation as the sequel to previous traumatic etiology of pancreatitis. The child has been planned for cystogastrostomy after 2 weeks of conservative management.

cystic mass of the organ. CFI further adds to the diagnosis to differentiate it from the vascular pathologies. MRI helps in anatomic details and further delineation of the

cystic mass. MRCP is an invaluable tool in finding any communication with the biliary or pancreatic tree. Conservative treatment is always preferred in children. This



includes bowel rest and total parenteral nutrition. The cutoff line for management is the size of the cyst. If the size is more than 5 cm, surgical maneuver is indicated, otherwise it is treated conservatively [3]. These cysts can become complicated by rupture, hemorrhage, or infection. Percutaneous drainage can be done safely, but recurrence is common. This is indicated in infected and non-septated pseudocysts. There was a complete resolution of pseudocysts in the series of 9 cases by Sharma and Maharshi [5] and all had endoscopic drainage [4,5].

### Conclusion

The management of pediatric pseudocyst depends on the size of the cyst and the associated extent of injury. The management can be contemplated by percutaneous, endoscopic procedure or internal drainage, depending on the guidelines described. Percutaneous drainage has always shown poorer outcome than surgical cases.

### Acknowledgement

We are thankful to Mr. Nitish Virmani, Ms Nitika, Rajdeep, and Raushan of the Faculty of Allied Health Sciences of SGT University, Gurgaon, for carrying out the radiological investigations and for procuring the images.

### List of Abbreviations

BTFE	Balanced turbo-field-echo
CECT	Contrast enhanced computerized tomography
CFI	Color flow imaging
CT	Computed Tomography
GB	Gall bladder
MRCP	Magnetic resonance cholangio-pancreaticography
MRI	Magnetic resonance imaging

### Summary of the case

<b>Patient (gender, age)</b>	1	3-year-old female child
<b>Final Diagnosis</b>	2	Post-traumatic pancreatic pseudocyst
<b>Symptoms</b>	3	Pain in abdomen following blunt trauma
<b>Medications</b>	4	Conservative treatment
<b>Clinical Procedure</b>	5	Underwent US, CECT, and MRI studies
<b>Specialty</b>	6	Radio-diagnosis

### Informed Consent

Informed consent of parents was taken to report this case.

### Competing Interest

None

### Author details

Shashi Sharma<sup>1</sup>, Bharat Bhushan Sharma<sup>2</sup>, Mir Rizwan Aziz<sup>2</sup>, Priyanka Rana<sup>2</sup>, Naveen Bhardwaj<sup>2</sup>

1. Department of Pediatrics, SGT Medical College, Budhera, Gurgaon, India.

2. Department of Radio-Diagnosis, SGT Medical College, Budhera, Gurgaon, India.

### References

1. Benifla M, Weizman Z. Acute pancreatitis in childhood: analysis of literature data. *J Clin Gastroenterol* 2003; 37(2):169–72. <https://doi:10.1097/00004836-200308000-00015>
2. Cabrera R, Otero H, Blesa E, Jimenez C, Nunez R. Pancreatic pseudocyst. Review of 22 cases. *Cir Pediatr* 1997; 10(2):49–53.
3. Shilyansky J, Sena LM, Kreller M, Chait P, Babyn PS, Filler RM, et al. Nonoperative management of pancreatic injuries in children. *J Pediatr Surg* 1998; 33(2):343–9. [https://doi:10.1016/S0022-3468\(98\)90459-6](https://doi:10.1016/S0022-3468(98)90459-6)
4. Teh SH, Pham TH, Lee A, Stavlo PL, Hanna AM, Moir C. Pancreatic pseudocyst in children: the impact of management strategies on outcome. *J Pediatr Surg* 2006; 41(11):1889–93. <https://doi:10.1016/j.jpedsurg.2006.06.017>
5. Sharma SS, Maharshi S. Endoscopic management of pancreatic pseudocyst in children: a long-term follow-up. *J Pediatr Surg* 2008; 43(9):1636–9. <https://doi:10.1016/j.jpedsurg.2008.01.026>