

# “Antibiotic-Responsive” drug fever due to the patient’s medication-taking behavior: a case report

Daiki Yokokawa<sup>1\*</sup> , Takanori Uehara<sup>1</sup> ,  
Yoshiyuki Ohira<sup>1,2</sup> , Kazutaka Noda<sup>1</sup> ,  
Masatomi Ikusaka<sup>1</sup> 

European Journal of Medical Case Reports

Volume 9(3):64–67

DOI: 10.24911/ejmcr.9-1624



This is an open access article distributed in accordance with the Creative Commons Attribution (CC BY 4.0) license: <https://creativecommons.org/licenses/by/4.0/> which permits any use, Share — copy and redistribute the material in any medium or format, Adapt — remix, transform, and build upon the material for any purpose, as long as the authors and the original source are properly cited. © The Author(s) 2025

## ABSTRACT

**Background:** Drug fever can complicate the diagnosis of recurrent fever, especially when multiple physicians are involved in treatment.

**Case Presentation:** We report the case of a 56-year-old man with recurrent fever responsive to antibiotics, initially suggesting a persistent bacterial infection. Detailed history-taking revealed that he discontinued minocycline, prescribed for rosacea dermatitis, when starting new antibiotics, and resumed it after the fever subsided. Eosinophilia and a clear improvement in the general condition between fevers led to the suspicion of drug fever. A rechallenge confirmed minocycline-induced fever.

**Conclusion:** This case emphasizes the importance of thoroughly reviewing medication history and patients’ medication-taking behavior when drug fever is suspected.

**Keywords:** Minocycline, drug fever, medication-taking behavior.

Received: 31 March 2025

Accepted: 10 December 2024

Type of Article: CASE REPORT

Specialty: Internal Medicine

Correspondence to: Daiki Yokokawa

\*Department of General Medicine, Chiba University Hospital, Chiba, Japan.

Email: dyokokawa6@gmail.com

Full list of author information is available at the end of the article.

## Background

The diagnosis of sustained fever can be complicated by medication-taking behavior, especially when medications are prescribed by multiple physicians. This report describes a case that was initially suspected to be “antibiotic-responsive” but was ultimately identified as minocycline-induced drug fever.

Factors contributing to poor medication adherence include patient-, physician-, and healthcare system-related factors [1]. Only approximately 50% of patients adhere to their prescribed medications, and care provided by multiple physicians affects medication adherence [1,2]. For example, a systematic review and meta-analysis involving 27 million patients with hypertension between 2010 and 2020 reported that 27%-40% exhibited poor adherence [3]. Additionally, a report conducted in Japan found that 24.7% of patients had discontinued prescribed antibiotics at their own discretion [4].

## Case Presentation

A 56-year-old Japanese male patient with recurrent fever was referred by his primary care physician (PCP) to our hospital. He developed a 39°C fever 19 days before the consultation and visited his PCP the next day. The physician prescribed levofloxacin 500 mg/day for 7 days, and the fever subsided after 4 days. However, the fever

recurred along with a cough 9 days before the consultation; hence, the PCP prescribed clarithromycin 400 mg/day for 5 days, resulting in fever reduction. While afebrile, the patient had a good appetite and no weight loss. Nevertheless, 3 days before the consultation, he developed a 38.9°C fever and was referred to our hospital. His medical history included atopic dermatitis, for which he had been seeing a dermatologist since the age of 25 years and was being treated with topical tacrolimus ointment and steroids. Additionally, he began taking minocycline for rosacea dermatitis 31 days before the consultation. The patient had no other medical history or history of medication use. There were no known allergies and no previous adverse drug reactions.

On physical examination, the patient was afebrile and had normal vital signs. In the head and neck examination, erythema without tenderness was noted on both cheeks, consistent with rosacea dermatitis. No conjunctival hemorrhage or jaundice was observed. The auricles were not deformed, and the oral cavity and pharynx appeared normal. The thyroid was normally palpable without tenderness. No cardiac murmurs or adventitious breath sounds were detected in the chest. The abdomen was non-tender with no hepatosplenomegaly. No joint swelling or muscle tenderness was noted in the extremities. There was no superficial lymphadenopathy. No abnormalities in the

skin were found other than the face. Blood tests revealed eosinophilia (2030/mcl) and an elevated C-reactive protein level (5.3 mg/dl). No anomalies suggesting neutrophilia, anemia, liver dysfunction, or renal impairment were found. Chest radiography results were normal.

The patient’s fever temporarily subsided with antibiotic treatment, raising suspicion of a persistent bacterial infection such as infectious endocarditis. A differential diagnosis was considered, including infectious endocarditis, pneumonia, and occult abscesses (Table 1). However, a review of systems revealed no abnormalities in vital signs such as tachypnea and tachycardia, and no findings suggestive of infective endocarditis, including heart murmurs, conjunctival hemorrhage, Janeway lesions, or Osler nodes. Moreover, his excellent general condition during the intermittent afebrile phases was atypical for a persistent bacterial infection, malignancy, and non-infectious inflammatory disease. In addition, the presence of eosinophilia was inconsistent with a typical bacterial

etiology. Detailed inquiries revealed the patient decided not to combine the dermatologist- and PCP-prescribed antibiotics. When he started the newly prescribed antibiotics, he discontinued minocycline; however, he resumed it after the fever subsided (Figure 1). Hence, we suspected minocycline-induced drug fever and conducted a rechallenge, which resulted in a high fever with chills after 4 hours. Subsequently, minocycline was discontinued, and no fever recurrence was observed.


**Discussion**

We report a case of recurrent fever with an antibiotic-response-like time course following antibiotic administration, which initially led to the consideration of persistent bacterial infections such as infective endocarditis, tuberculosis, and occult abscesses. However, the diagnosis was later reconsidered in light of the patient’s preserved general condition during afebrile periods. A detailed history revealed that the patient discontinued minocycline on his own when levofloxacin and clarithromycin were prescribed. He had been intermittently discontinuing and restarting minocycline, which better explained the observed clinical pattern. The patient’s condition was not a medication effect, but rather a side effect.

When considering the causes of fever of unknown origin, factors such as infection, connective tissue diseases, and malignancy are commonly considered. However, when the patient’s general condition clearly improves during afebrile periods, drug-induced fever should be considered in the differential diagnosis. A 2023 review defines drug fever as a febrile response that meets all the following criteria: (i) onset after drug administration, (ii) resolution within 72 hours after discontinuation of the drug without specific treatment, (iii) no other identifiable cause based on medical history, physical examination, laboratory tests, or imaging, and (iv) no recurrence of fever within 72 hours after defervescence [6]. A retrospective

**Table 1.** A list of common causes of sustained fever (modified from the reference [5]).

|   |  |
|---|--|
| <b>INFECTION</b>                          | Bacterial<br>Sinusitis<br>Dental abscesses<br>Endocarditis<br>Tuberculosis<br>Abdominal or pelvic abscess<br>Urinary tract infection<br>Viral<br>Cytomegalovirus<br>Epstein-Barr virus |
| <b>MALIGNANCY</b>                         | Leukemia<br>Lymphoma   |
| <b>NONINFECTIOUS INFLAMMATORY DISEASE</b> | Connective tissue diseases<br>Granulomatous disease<br>Vasculitis  |
| <b>MISCELLANEOUS</b>                      | Drug-induced<br>Factitious fever<br>Thromboembolic disease<br>Thyroiditis  |



**Figure 1.** The course of symptoms and medications.

study at a single institution in Japan reported that drug fever accounted for 5.7% of hospitalized patients presenting with fever [7].

Whether intentionally or unintentionally, patients may not report their exact medication usage; therefore, medical staff should specifically ask about their use of medicines and other treatments to rule out drug-induced fever. It is important to ensure that all medications taken within the past month, including those prescribed by other physicians and over-the-counter drugs, are accounted for [8]. Patients often cannot accurately report their medication history and may not bring either the medications or a recent list of them [9]. Utilizing pharmacy records from community pharmacists may provide more comprehensive information regarding the medication history of hospitalized patients [9].

Initially, the drug-induced fever developed approximately 13 days after the minocycline prescription, and the interval between administration and fever onset progressively shortened. This suggests an immune response as the underlying mechanism of drug-induced fever [10]. Drug hypersensitivity elicits different immune responses depending upon the nature of the drug and varies between individuals. For instance, drug-induced immune hemolytic anemia are type II hypersensitivity reactions, whereas, drug-induced hypersensitivity syndrome, is a type IV hypersensitivity reaction. Drug fever may be categorized as a type III reaction among the four classical types of allergic responses [11]. In patients with serum sickness, a condition similar in nature to drug fever, decreased levels of serum C3 and C4 have been observed, along with immune complex deposition in the vasculature, supporting its classification as a type III reaction. However, drug fever may also elicit type IV or delayed type hypersensitivity reaction, as seen in the case of drug fever induced by Vinca alkaloids [12]. Even if the patient was asymptomatic during the prior administration, sensitization may have occurred, and re-exposure can trigger symptoms. In addition, prior exposure to a drug of the same class has been associated with a shortened latency period before the onset of drug fever [13]. Research on whether atopic dermatitis constitutes a risk factor for drug fever remains insufficient. A pediatric study conducted in China reported that atopic dermatitis posed a risk for delayed-type drug hypersensitivity, with an odds ratio of 8.2 [14]. However, another report found that atopic dermatitis was not a significant risk factor for drug hypersensitivity induced by antimicrobial agents [9]. When drug-induced fever is suspected in mild cases, a rechallenge can be safely performed to facilitate diagnosis. However, rechallenges are contraindicated in patients with severe liver injury, lymph node swelling, or suspected drug-induced hypersensitivity syndrome.

## Conclusion

This case underscores the importance of considering drug-induced fever in the differential diagnosis of recurrent fever, especially when eosinophilia is present. A detailed medication history and awareness of patient medication-taking behaviors are essential to avoid unnecessary antibiotic use and misdiagnosis of persistent bacterial infections.

### What is new

This case underscores the importance of considering drug-induced fever in the differential diagnosis of recurrent fever, especially when eosinophilia is present. A detailed medication history and awareness of patient medication-taking behaviors are essential to avoid unnecessary antibiotic use and misdiagnosis of persistent bacterial infections.

## Acknowledgment

None.

## List of abbreviations

PCP Primary care physician

## Conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this article.

## Consent for publication

Written informed consent was obtained from the patient.

## Funding

None.

## Ethical approval

Ethical approval is not required at our institution for anonymous case reports.

## Author details

Daiki Yokokawa<sup>1</sup>, Takanori Uehara<sup>1</sup>, Yoshiyuki Ohira<sup>1,2</sup>, Kazutaka Noda<sup>1</sup>, Masatomi Ikusaka<sup>1</sup>

1. Department of General Medicine, Chiba University Hospital, Chiba, Japan
2. Department of General Internal Medicine, St. Marianna University School of Medicine, Kawasaki, Japan

## References

1. Brown MT, Bussell JK. Medication adherence: WHO cares? *Mayo Clin Proc*. 2011 Apr;86(4):304–14. <https://doi.org/10.4065/mcp.2010.0575>
2. Sabaté E, editor. *Adherence to long-term therapies: evidence for action*. Geneva, Switzerland: World Health Organization; 2003.
3. Lee EK, Poon P, Yip BH, Bo Y, Zhu MT, Yu CP, et al. Global burden, regional differences, trends, and health consequences of medication nonadherence for hypertension during 2010 to 2020: a meta-analysis involving 27 million patients. *J Am Heart Assoc*. 2022 Sep;11(17):e026582. <https://doi.org/10.1161/JAHA.122.026582>
4. National Center for Global Health and Medicine, Antimicrobial Resistance Clinical Reference Center. *Antimicrobial drug awareness survey report 2024*. 2024

- [cited 2025 Feb 26];11. Available from: [https://amr.ncgm.go.jp/pdf/20241004\\_report.pdf](https://amr.ncgm.go.jp/pdf/20241004_report.pdf)
5. Hersch EC, Oh RC. Prolonged febrile illness and fever of unknown origin in adults. *Am Fam Physician*. 2014 Jul;90(2):91–6.
  6. Someko H, Kataoka Y, Obara T. Drug fever: a narrative review. *Ann Clin Epidemiol*. 2023 May;5(4):95–106. <https://doi.org/10.37737/ace.23013>
  7. Ryuko H, Otsuka F. A comprehensive analysis of 174 febrile patients admitted to Okayama University Hospital. *Acta Med Okayama*. 2013;67(4):227–37.
  8. Fitzgerald RJ. Medication errors: the importance of an accurate drug history. *Br J Clin Pharmacol*. 2009 Jun;67(6):671–5. <https://doi.org/10.1111/j.1365-2125.2009.03424.x>
  9. Kühl J, Bergh B, Laudes M, Szymczak S, Heine G. Cofactors of drug hypersensitivity-A monocenter retrospective analysis. *Front Allergy*. 2023 Jan;3:1097977. <https://doi.org/10.3389/falgy.2022.1097977>
  10. Mackowiak PA. Drug fever: mechanisms, maxims and misconceptions. *Am J Med Sci*. 1987 Oct;294(4):275–86. <https://doi.org/10.1097/00000441-198710000-00011>
  11. Lawley TJ, Bielory L, Gascon P, Yancey KB, Young NS, Frank MM. A prospective clinical and immunologic analysis of patients with serum sickness. *N Engl J Med*. 1984 Nov;311(22):1407–13. <https://doi.org/10.1056/NEJM198411293112204>
  12. Kidon MI, Haj Yahia S, Abebe-Campino G, Agmon-Levin N, Yelon M. Drug fever-an immune-mediated delayed type hypersensitivity reaction to Vinca alkaloids in pediatric oncology patients, possibly mediated by cysteinyl leukotrienes. *Front Allergy*. 2024 Aug;5:1361403. <https://doi.org/10.3389/falgy.2024.1361403>
  13. Oizumi K, Onuma K, Watanabe A, Motomiya M. Clinical study of drug fever induced by parenteral administration of antibiotics. *Tohoku J Exp Med*. 1989 Sep;159(1):45–56. <https://doi.org/10.1620/tjem.159.45>
  14. Duque JS, Cheuk DK, Chong PC, Ip P, Wong WH, Lee PP, et al. Risk factors for drug allergies in Chinese children. *Asian Pac J Allergy Immunol*. 2020 Dec;38(4):271–8.

### Summary of case

|   |                       |                                |
|---|-----------------------|--------------------------------|
| 1 | Patient (gender, age) | 56 years, male                 |
| 2 | Final diagnosis       | Minocycline-induced drug fever |
| 3 | Symptoms              | Recurrent fever                |
| 4 | Medications           | NA                             |
| 5 | Clinical procedure    | Minocycline was discontinued   |
| 6 | Specialty             | Internal medicine              |