

68 **Clinical examination**

69 Two contiguous nodular fusiform swellings measuring
70 approximately 14 × 4 cm were present along the flexor
71 aspect of the forearm, extending from the antecubital
72 fossa to the wrist and continuing to the palmar aspect
73 of the ring finger. The masses were firm, mildly tender,
74 and showed restricted mobility along the longitudinal
75 axis. Percussion over the swelling elicited tingling in the

median nerve distribution (positive Tinel’s sign). Radial 76
and ulnar arteries were palpable. No cutaneous stigmata 77
of NF were observed. 78

Diagnostic assessment 79

Imaging 80

MRI demonstrated a well-defined, elongated, multi- 81
lobulated lesion along the median nerve involving the 82




Figure 1. MRI of the right forearm demonstrating a well-defined, elongated, multilobulated lesion along the course of the median nerve within the flexor compartment.




Figure 2. MRI of the wrist and hand showing distal extension of the lesion along the median nerve into the palmar aspect of the hand.

83 flexor compartment of the forearm with extension to
84 the wrist and palm. The lesion appeared hypointense on
85 T1-weighted images and hyperintense on T2-weighted
86 images with areas of cystic degeneration (Figures 1
87 and 2).

Histopathological examination

Core needle biopsy revealed spindle cells arranged in palisading patterns. Immunohistochemistry (IHC) demonstrated strong **S-100 positivity** with a Ki-67 index <5%, confirming schwannoma.

88

89

90

91

92



Figure 3. Intraoperative photograph demonstrating exposure of the tumor beneath the flexor tendons within the flexor compartment of the forearm.




Figure 4. Intraoperative view showing the tumor arising eccentrically from the median nerve with displacement of surrounding nerve fascicles.

93 **Treatment and intervention**

94 **Surgical procedure**

95 The patient underwent complete surgical excision of the
96 tumor with preservation of the median nerve. The opera-
97 tion involved careful dissection under 2.5× magnification

98 using a surgical loupe. A longitudinal incision had been 98
99 centered over the tumor bulk, extending from the elbow 99
100 to the wrist. The incision was deepened to the deep fascia. 100
101 A plane was created between the flexor carpi radialis and 101
102 the palmaris longus muscles (Figure 3). The median nerve 102
103 was identified below the radial attachment of the flexor 103



Figure 5. Intraoperative dissection of the wrist and palm demonstrating extension of the tumor along the median nerve with preservation of adjacent neurovascular structures.




Figure 6. Final intraoperative view after tumor excision showing intact median nerve branches with preservation of neural continuity.

104 digitorum superficialis. The tumor was firmly adherent to
105 the median nerve and had an eccentric location (Figure
106 4). Using fine scissors, the soft tumor mass was separated
107 from the surrounding nerve fascicles as well as median
108 nerve branches (anterior interosseous nerve as well as

109 muscle branches) after the epineurium was cut longitu-
110 dinally (Figure 5). The incision was later extended over
111 the wrist. Meticulous dissection was performed without
112 the use of a tourniquet. We believe that identifying min-
113 ute vessels is better done this way, which is important for




Figure 7. Gross specimen of the excised tumor demonstrating encapsulated nodular schwannomas removed from the median nerve.




Figure 8. Pre- and post-op histopathological images. (A). Low-power scanner view ($\times 4$) demonstrating an encapsulated, well-circumscribed tumor composed of spindle cells arranged in intersecting fascicles (upper left). (B). Immunohistochemical staining showing a low Ki-67 proliferation index ($\sim 5\%$) despite the presence of occasional bizarre nuclei (upper middle). (C). IHC demonstrating diffuse S-100 positivity in the tumor cells, supporting the diagnosis of schwannoma (upper right). (D). High-power view ($\times 40$) demonstrating areas of ancient change characterized by scattered atypical and bizarre nuclei (blue arrows), representing degenerative changes (lower left). (E). Low-power view ($\times 10$) showing intersecting fascicles of spindle cells with formation of a Verocay body (blue arrow), characterized by nuclear palisading and an intervening anuclear fibrillary zone (lower middle). (F). Low-power view ($\times 10$) showing numerous hyalinized vessels within the tumor, a characteristic feature frequently observed in schwannomas (lower right).

114 preserving vasa nervosa. Staged incisions were made on
 115 the palmar aspect of the hand, extending to the ring finger
 116 in a zig-zag pattern. The flexor retinaculum was cut with a
 117 blade, and the median nerve was identified as it travelled
 118 posteriorly toward the middle finger as well as the lateral
 119 aspect of the ring finger. Posteriorly, below the flexor ret-
 120 inaculum's attachment on the wrist's medial aspect, the
 121 ulnar nerve, as well as the ulnar artery, were identified
 122 in Guyon's canal and preserved. The tumor did not arise
 123 from the ulnar nerve fibers, although its extension was
 124 toward the medial aspect of the ring finger. The tumor
 125 was completely removed, preserving all nerve fibers to
 126 the ring finger (Figures 6 and 7). The wound was closed
 127 in layers, and a 14 Fr suction drain was placed, which was
 128 removed on postoperative day 1.

129 *Outcome*

130 In the immediate postoperative period, on-table assess-
 131 ment following reversal of neuromuscular blockade with
 132 vecuronium demonstrated an optimal functional outcome,
 133 with improvement in the preoperative deficits likely
 134 caused by nerve compression (Supplementary 1 video).
 135 Video link: <https://youtube.com/shorts/fM07IHBpFpI>

136 *Final histopathology report*

137 Grossly, there was an encapsulated white-grey soft tis-
 138 sue mass with yellow areas. Microscopically, the sec-
 139 tion shows an encapsulated tumor composed of compact
 140 hypercellular areas of spindle cell fascicles with ill-de-
 141 fined cytoplasm. Occasional areas of ancient change with
 142 atypical cells and blood vessels with thickened hyalinized
 143 walls were noted. This suggests a cellular schwannoma
 144 with focal ancient change (Figure 8).

145 *Postoperative care*

146 The patient was discharged on postoperative day 1. She
 147 received physical rehabilitation on an outpatient basis in
 148 our dedicated oncorehab unit.

149 *Follow-up and outcomes*

150 *Short-term outcomes*

151 A short-term follow-up was conducted 2 weeks after sur-
 152 gery, during which staplers and sutures were removed.
 153 Sensory neural functions were not impaired.

154 Long-term follow-up is planned with periodic clinical
 155 examination to monitor for recurrence and assess neu-
 156 rological function. Although recurrence after complete
 157 excision of a schwannoma is uncommon, continued sur-
 158 veillance is recommended.

159 **Discussion**

160 Schwannomas are benign peripheral nerve sheath tumors
 161 that typically occur as solitary lesions. Extensive involve-
 162 ment of the median nerve by multiple schwannomas

spanning several anatomical compartments is exception- 163
 ally rare and poses unique diagnostic and surgical chal- 164
 lenges [4,5]. 165

The present case demonstrates an unusually long-seg- 166
 ment involvement of the median nerve extending from the 167
 elbow to the distal interphalangeal joint. Most reported 168
 cases describe isolated schwannomas located at the wrist 169
 or forearm, whereas extensive lesions involving multi- 170
 ple compartments of the upper limb are rarely described 171
 [6]. Although the presence of multiple schwannomas has 172
 historically been referred to as schwannomatosis, cur- 173
 rent classification systems define schwannomatosis as a 174
 distinct entity within the NF spectrum requiring specific 175
 clinical or genetic criteria. In our patient, the lesions were 176
 confined to the median nerve without evidence of sys- 177
 temic disease or genetic confirmation; she does not fulfil 178
 the updated 2022 criteria for NF-2-related schwannom- 179
 atosis as described by Plotkin et al. [7] Therefore, this 180
 presentation is more appropriately interpreted as multiple 181
 schwannomas arising along the median nerve rather than 182
 true schwannomatosis. 183

Preoperative diagnosis of peripheral nerve sheath tum- 184
 ors can be challenging. Clinically, these lesions may present 185
 as slowly enlarging masses associated with pain, pares- 186
 thesia, or a positive Tinel's sign. Imaging modalities such 187
 as MRI are useful for demonstrating tumor extent and its 188
 relationship to adjacent structures, although distinguishing 189
 schwannomas from neurofibromas solely on imaging may 190
 be difficult [8]. Histologically, schwannomas are charac- 191
 terized by a well-defined capsule derived from epineurium 192
 and demonstrate alternating Antoni A and Antoni B pat- 193
 terns, with strong S-100 immunoreactivity [9]. 194

From a surgical perspective, schwannomas can usually 195
 be excised while preserving the parent nerve because they 196
 arise eccentrically from the nerve sheath. Careful identi- 197
 fication of proximal and distal nerve segments, magnified 198
 dissection, and meticulous separation of the tumor cap- 199
 sule from the splayed nerve fascicles are essential steps to 200
 avoid neurological injury. Recurrence following complete 201
 excision is uncommon, and most patients experience sig- 202
 nificant symptomatic improvement with preservation of 203
 nerve function. Das et al. [10] reported improvement of 204
 preoperative symptoms with maintained nerve function in 205
 approximately 89% of patients undergoing surgical exci- 206
 sion of schwannomas. 207

208 **Conclusion**

209 Complete surgical excision with preservation of the parent 210
 nerve remains the treatment of choice for median nerve 211
 schwannomas. Careful microsurgical dissection allows 212
 tumor removal while maintaining neural integrity and 213
 functional outcome. Even in cases with extensive involve- 214
 ment of the median nerve, nerve-preserving surgery can 215
 achieve excellent symptomatic and functional recovery.

217 **Supplementary Material**
 218 Video available here:
 219 https://youtube.com/shorts/xWtiu_vde9w

220 **What's new?**
 221 The authors describe an unusual case of extensive multiple
 222 schwannomas involving the median nerve, extending from
 223 the elbow to the distal interphalangeal joint across several
 224 anatomical compartments. The case highlights the diagnos-
 225 tic challenges, the role of preoperative tissue diagnosis, and
 226 the feasibility of meticulous nerve-preserving surgical exci-
 227 sion resulting in excellent functional recovery.

228 **List of Abbreviations**
 229 IHC Immunohistochemistry
 230 MRI Magnetic resonance imaging
 231 NF Neurofibromatosis

232 **Conflict of interest**
 233 The authors declare that they have no conflicts of interest.

234 **Funding**
 235 This research did not receive any specific grants from funding
 236 agencies in the public, commercial, or not-for-profit sectors.

237 **Consent for publication**
 238 Written informed consent was obtained from the patient for
 239 publication of this case report and any accompanying images.

240 **Ethical approval**

241 **Patient perspective**
 242 The patient reported significant relief of symptoms following
 243 surgery and was satisfied with the cosmetic and functional
 244 outcomes.

245 **Data availability**
 246 Data will be available on request due to privacy/ethical restric-
 247 tions. The data that support the findings of this study are availa-
 248 ble on request from the corresponding author. The data are not
 249 publicly available due to privacy or ethical restrictions.

250 **Author details**
 251 Mohit Malhotra¹, Monmeeta Srikant², Tushar Deka¹, Lopamudra
 252 Kakoti², Gaurav Das¹
 253 1. Department of Surgical Oncology, Dr. B. Borooah Cancer
 254 Institute, Guwahati, India

2. Department of Oncopathology, Dr. B. Borooah Cancer 255
 Institute, Guwahati, India 256

References 257

1. Majumder A, Ahuja A, Chauhan D, Paliwal P, Bhardwaj 258
 M. A clinicopathological study of peripheral schwanno- 259
 mas. *Med Pharm Rep*. 2021;94(2):191–6. [https://doi.](https://doi.org/10.15386/mpr-1708) 260
[org/10.15386/mpr-1708](https://doi.org/10.15386/mpr-1708) 261
2. Aydin M, Kotan D, Keles M. Acute median nerve palsy due 262
 to hemorrhaged schwannoma: case report. *J Brachial Plex* 263
Peripher Nerve Inj. 2007;2:19. 264
3. Kütahya H, Güleç A, Güzel Y, Kacira B, Toker S. 265
 Schwannoma of the median nerve at the wrist and 266
 palmar regions of the hand: a rare case report. *Case* 267
Rep Orthop. 2013;2013(1):950106. [https://doi.](https://doi.org/10.1155/2013/950106) 268
[org/10.1155/2013/950106](https://doi.org/10.1155/2013/950106) 269
4. Padasali P, Shankaregowda V, Kshirsagar S. Median 270
 nerve schwannoma: a case and review of literature. 271
Asian J Neurosurg. 2015;10(3):212–5. [https://doi.](https://doi.org/10.4103/1793-5482.161178) 272
[org/10.4103/1793-5482.161178](https://doi.org/10.4103/1793-5482.161178) 273
5. Hanemann CO, Evans DG. News on the genetics, epidemi- 274
 ology, medical care and translational research of schwan- 275
 nomas. *J Neurol*. 2006;253(12):1533–41. [https://doi.](https://doi.org/10.1007/s00415-006-0347-0) 276
[org/10.1007/s00415-006-0347-0](https://doi.org/10.1007/s00415-006-0347-0) 277
6. Gonzalvo A, Fowler A, Cook RJ, Little NS, Wheeler H, 278
 McDonald KL, et al. Schwannomatosis, sporadic schwan- 279
 nomatosis, and familial schwannomatosis: a surgical series 280
 with long-term follow-up. *J Neurosurg*. 2011;114(3):756– 281
 62. <https://doi.org/10.3171/2010.8.JNS091900> 282
7. Plotkin SR, Messiaen L, Legius E, Pancza P, Avery RA, 283
 Blakeley JO, et al. Updated diagnostic criteria and nomen- 284
 clature for neurofibromatosis type 2 and schwannomato- 285
 sis: an international consensus recommendation. *Genet* 286
Med. 2022;24(9):1967–77. [https://doi.org/10.1212/](https://doi.org/10.1212/WNL.98.18_supplement.3996) 287
[WNL.98.18_supplement.3996](https://doi.org/10.1212/WNL.98.18_supplement.3996) 288
8. Pilavaki M, Chourmouzi D, Kiziridou A, Skordalaki A, 289
 Zarampoukas T, Drevelengas A. Imaging of peripheral 290
 nerve sheath tumors with pathologic correlation: picto- 291
 rial review. *Eur J Radiol*. 2004;52(3):229–39. [https://doi.](https://doi.org/10.1016/j.ejrad.2003.12.001) 292
[org/10.1016/j.ejrad.2003.12.001](https://doi.org/10.1016/j.ejrad.2003.12.001) 293
9. Pekmezci M, Reuss DE, Hirbe AC, Dahiya S, Gutmann 294
 DH, Von Deimling A, et al. Morphologic and immu- 295
 nohistochemical features of malignant peripheral 296
 nerve sheath tumors and cellular schwannomas. *Mod* 297
Pathol. 2015;28(2):187–200. [https://doi.org/10.1038/](https://doi.org/10.1038/modpathol.2014.109) 298
[modpathol.2014.109](https://doi.org/10.1038/modpathol.2014.109) 299
10. Das S, Ganju A, Tiel RL, Kline DG. Tumors of the brachial 300
 plexus. *Neurosurg Focus*. 2007;22(6):E26. [https://doi.](https://doi.org/10.3171/foc.2007.22.6.27) 301
[org/10.3171/foc.2007.22.6.27](https://doi.org/10.3171/foc.2007.22.6.27) 302

Summary of case

1	PATIENT (GENDER, AGE)	17-year-old female
2	FINAL DIAGNOSIS	Schwannoma
3	SYMPTOMS	Numbness and paraesthesia
4	MEDICATIONS	None (managed surgically)
5	CLINICAL PROCEDURE	Surgical excision of the mass
6	SPECIALTY	Oncology