Unilateral Graves' disease

Rodrigo Sanchez Magallan Gonzalez[1]*, Mercedes Aguilar Soto[2]

European Journal of Medical Case Reports
Volume XX(XX):XX–XX
DOI: XXXX

OPEN ACCESS

This is an open access article distributed in accordance with the Creative Commons Attribution (CC BY 4.0) license: https://creativecommons.org/licenses/by/4.0/) which permits any use, Share — copy and redistribute the material in any medium or format, Adapt — remix, transform, and build upon the material for any purpose, as long as the authors and the original source are properly cited. © The Author(s) 2026

Type of Article: Image **Specialty:** Endocrinology

Correspondence to: Rodrigo Sanchez Magallan Gonzalez, MD
*Department of Internal Medicine, The American British Cowdray Medical Center. Av. Carlos Graef Fernández 154, Santa Fe, Cuajimalpa de Morelos, 05300 Mexico City, Mexico.
Email: dr.rodrigosmg@gmail.com
Full list of author information is available at the end of the article.

Received: 15 September 2026
Revised (1): 13 October 2026
Accepted: 16 October 2026

Learning Points

1. Graves' disease can present with tracer uptake confined to a single thyroid lobe; this unilateral variant accounts for fewer than 30 reported cases and is not addressed in current guidelines.
2. Unilateral uptake on scintigraphy is not diagnostic on its own: cervical ultrasonography is required to demonstrate a structurally present contralateral lobe and to exclude thyroid hemiagenesis and a hyperfunctioning nodule, and thyrotropin receptor antibodies are required to establish autoimmune aetiology.
3. The apparently spared lobe is suppressed rather than absent and remains capable of autoimmune stimulation, so hyperthyroidism may recur in it after hemithyroidectomy; total thyroidectomy is therefore worth considering should surgery be chosen; the published experience is too limited to establish it as a formal recommendation.

Case Description

A 45-year-old man presented to the emergency department with dyspnea and chest pain. Electrocardiography showed sinus tachycardia, and cardiac biomarkers were negative. He reported increased stool frequency over the preceding month and unintentional weight loss of 5 kg over two months, without fever, night sweats, or lymphadenopathy.

Thyroid function testing showed a suppressed thyrotropin of 0.0009 mIU/L (reference range 0.4 to 4.0 mIU/L) and an elevated free thyroxine of 5.6 ng/dL (reference range 0.8 to 1.8 ng/dL). Thyrotropin receptor antibodies were positive. Technetium-99m pertechnetate scintigraphy (Figure 1) demonstrated intense, homogeneous tracer accumulation confined to the right lobe, with only faint activity over the isthmus and left lobe. Cervical ultrasonography confirmed a normally located left lobe of preserved volume, excluding thyroid hemiagenesis, and showed no nodule in either lobe. Together with the positive thyrotropin receptor antibodies, these findings established the diagnosis of unilateral Graves' disease. The patient was started on propranolol and thiamazole (methimazole); at 5 months, he was biochemically euthyroid.

Discussion

Unilateral Graves' disease (autoimmune hyperthyroidism in which tracer uptake is confined to one lobe of an anatomically bilobar gland) is rare: a systematic review published in 2025 identified only 27 reported patients, the right lobe being involved in roughly two-thirds of them [1]. The scan alone points away from the diagnosis because markedly unilateral uptake is more often produced by a toxic adenoma, by thyroid hemiagenesis with Graves' disease in the remaining lobe, or by unilateral thyroiditis [1,2]. The diagnosis therefore rests on the combination of cervical ultrasonography demonstrating two structurally normal lobes without a nodule and positive thyrotropin receptor antibodies [1,3]. The distinction matters because the apparently silent lobe is suppressed rather than absent and remains capable of autoimmune stimulation: contralateral recurrence after hemithyroidectomy has been reported, and although the published experience is too small to support a firm recommendation, total thyroidectomy is worth considering if surgery is chosen [1].

Keywords: Graves' disease, thyroid scintigraphy, technetium-99m pertechnetate, thyrotropin receptor antibodies, thyrotoxicosis, thyroid hemiagenesis.

Informed consent

Written informed consent for publication of this case and the accompanying image was obtained from the patient.




Figure 1. *Anterior technetium-99m pertechnetate thyroid scintigraphy. Intense, homogeneous tracer accumulation is confined to the right lobe; the isthmus and left lobe show only faint background-level activity. The study is displayed in the anterior projection, so the patient's right lobe appears on the left of the image. R, right; L, left.*

Conflicts of interest

The authors declare that he has no conflicts of interest.

Funding

This work received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author details

Rodrigo Sanchez Magallan Gonzalez, MD[1], Mercedes Aguilar Soto[2]

1. Department of Internal Medicine, The American British Cowdray Medical Center. Av. Carlos Graef Fernández 154, Santa Fe, Cuajimalpa de Morelos, 05300 Mexico City, Mexico
2. Department of Endocrinology, The American British Cowdray Medical Center. Av. Carlos Graef Fernández 154, Santa Fe, Cuajimalpa de Morelos, 05300 Mexico City, Mexico


References

1. Scappaticcio L, Caruso P, Longo M, Volatile A, Cirillo P, Di Maio F, et al. Unilateral Graves' disease: a case report with concomitant thyroid cancer and systematic review of literature. Minerva Surg. 2025;80(3):274–281. https://doi.org/10.23736/S2724-5691.25.10685-0
2. Shechner C, Kraiem Z, Zuckerman E, Dickstein G. Toxic Graves' disease with thyroid hemiagenesis: diagnosis using thyroid-stimulating immunoglobulin measurements. Thyroid. 1992;2(2):133–135. https://doi.org/10.1089/thy.1992.2.133
3. Ross DS, Burch HB, Cooper DS, Greenlee MC, Laurberg P, Maia AL, et al. 2016 American Thyroid Association guidelines for diagnosis and management of hyperthyroidism and other causes of thyrotoxicosis. Thyroid. 2016;26(10):1343–1421. https://doi.org/10.1089/thy.2016.0229


Additional thyroid scintigraphy images

These images were obtained from the same technetium-99m pertechnetate thyroid scintigraphy already reported in the manuscript.

No image has been altered other than by cropping and uniform resizing. Images are displayed in the anterior projection, so the patient's right lobe appears on the left of each panel. R, right; L, left.




Panel A. *Anterior view, rainbow (blue-background) colour scale. Tracer is seen in the salivary glands and nasal mucosa, with intense accumulation in the right thyroid lobe and minimal activity in the left lobe.*




Panel B. *Rainbow colour scale over a different background window.*



Panel C. *Inverted greyscale.*



Panel D. *Inverted hot-metal colour scale.*



Panel E. *The right lobe reaches the top of the scale while the left lobe and isthmus remain at very low intensity.*