

Figure 1. Polymorphic VT with varying morphology VPCs at the time of presentation to the emergency department.




Figure 2. Markedly prolonged QT interval (QTc—560 to 580 ms, Bazett formula) recorded on the ECG that was taken soon after cardioversion of VT.

58 Blood tests revealed a serum potassium level of 3
59 mmol/l, while all other parameters, including arterial
60 blood gases, were normal. Intravenous potassium correc-
61 tion was initiated, and a beta blocker (Metoprolol succi-
62 nate 25 mg once daily) was added to her treatment plan.
63 With these measures, there was no recurrence of VT. A

bedside echocardiogram revealed a structurally normal 64
heart. She had an uneventful recovery, and by the third 65
day, her QT interval had normalized to a corrected QT of 66
436 ms (Figure 3). 67
The patient disclosed that she had been consuming 68
a crushed ginseng root (Figure 4) mixture with water 69




Figure 3. Normal QT interval on day 3 of admission (QTc - 436 ms).



Figure 4. Dried Ginseng root consumed by the patient.

twice a day as a traditional remedy for hemorrhoids. She had been doing this intermittently for 2 months, but recently increased her intake to larger quantities (5-8 g of raw dried ginseng root daily). She was on absolute fasting (abstaining from both solid and liquid food intake) for 2 days before the cardiac emergency occurred, during which she continued to take the ginseng mixture. She is a mother of three and reported no significant family or past medical history. She was not on any other medication, and there was no history to suggest substance abuse.

Discussion

Although ginseng is a well-known medicinal herb, there have been reports of potentially serious cardiac adverse effects. A woman developed long QT with torsades de pointes after consuming ginseng daily for over 6 months [5]. Such adverse reactions are likely rare and considered idiosyncratic. Nonetheless, the effect of ginseng on the QT interval has been demonstrated in a randomized prospective study, where participants taking ginseng extracts for 28 days exhibited a significant prolongation of the QTc compared to the placebo group [6].

In a study involving animal subjects, the infusion of Rg1 ginsenosides (the bioactive component of ginseng) resulted in the prolongation of the ventricular effective refractory period [7]. Another bioactive component of ginseng, ginsenoside Re, has been shown to suppress the rapid component of the delayed rectifier potassium current (IKr) [8]. This current is crucial for repolarization of the heart muscle cells, and blocking IKr can prolong the ventricular action potential duration, causing delayed repolarization and a prolonged QT interval [9].

The patient also had a low serum potassium level of 3 mmol/l (Normal reference range 3.5 - 5 mmol/l). It is unclear if ginseng directly contributes to hypokalemia, and hence, the association between ginseng intake and reduced potassium levels remains speculative.

Dyselectrolytemia and prolonged QTc during fasting is a well-documented phenomenon [10,11]. In addition to reduced intake during fasting, there is rapid potassium loss through renal excretion [12]. The Volume loss associated with fasting triggers the secretion of aldosterone, which stimulates the reabsorption of sodium and the excretion of potassium, leading to hypokalemia. In addition, metabolic alkalosis observed during fasting leads to bicarbonate diuresis and loss of accompanying potassium ions in the urine [13]. Even though the duration of fasting in this particular case is short (2 days), the possibility that this was a contributing factor to the cardiac arrhythmia cannot be ignored.

Panax pseudoginseng is native to India and is particularly abundant in the Northeast. Despite its presence in various parts of the region, its survival is threatened by overexploitation for medicinal use [14]—chrome-extension://efaidnbmnnnibpcajpcgclcfndmkaj/https://ijcs.ro/public/IJCS-16-61_Jamir.pdf

Conclusion

Given the widespread use of ginseng, there is a risk of inadvertent misuse leading to potentially life-threatening adverse effects. Clinicians should actively inquire about herbal supplement use in patients with unexplained arrhythmias or QT prolongation. It is also essential to disseminate information about such adverse effects to the public.

What is new

Indiscriminate use of traditional medicinal herbs like ginseng can lead to serious and life-threatening cardiac arrhythmias.

List of Abbreviations

ECG	Electrocardiogram
ED	Emergency department
PVC	Premature ventricular complexes
VT	Ventricular tachycardia

Conflict of interests

The authors declare that there is no conflict of interest regarding the publication of this article.

Funding

None.

Consent for publication

Due permission was obtained from the patient to publish the case and the accompanying images.

Ethical approval

Ethical approval is not required at our institution to publish an anonymous case report.

Author details

Avitso Liegise¹, Lanu Naro², Pouveine David³, Gailuna Maringmei³, Krosakhol Pucho¹

1. Senior Consultant, Department of Cardiology, Christian Institute of Health Sciences and Research (CIHSR), Dimapur, India
2. Nursing Supervisor, Department of Cardiology, Christian Institute of Health Sciences and Research, Dimapur, India
3. Nurse, Department of Cardiology, Christian Institute of Health Sciences and Research, Dimapur, India

References

1. Potenza MA, Montagnani M, Santacroce L, Charitos IA, Bottalico L. Ancient herbal therapy: a brief history of *Panax ginseng*. *J Ginseng Res*. 2023; 47(3):359–65.; <https://doi.org/10.1016/j.jgr.2022.03.004>
2. Yang WZ, Hu Y, Wu WY, Ye M, Guo DA. Saponins in the genus *Panax* L. (*Araliaceae*): a systematic review of their chemical diversity. *Phytochemistry*. 2014; 106:7–24.; <https://doi.org/10.1016/j.phytochem.2014.07.012>
3. Aminifard T, Razavi BM, Hosseinzadeh H. The effects of ginseng on the metabolic syndrome: an updated review. *Food Sci Nutr*. 2021; 9(9):5293–311.; <https://doi.org/10.1002/fsn3.2475>
4. Kim JW, Han SW, Cho JY, Chung JJ, Kim JG, Lee KH, et al. Korean red ginseng for cancer-related fatigue in colorectal cancer patients with chemotherapy: a randomised phase III trial. *Eur J Cancer*. 2020; 130:51–62.; <https://doi.org/10.1016/j.ejca.2020.02.018>
5. Torbey E, Abi Rafeh N, Khoueiry G, Kowalski M, Bekheit S. Ginseng: a potential cause of long QT. *J Electrocardiol*. 2011; 44(3):357–8.; <https://doi.org/10.1016/j.jelectrocard.2010.08.007>
6. Caron MF, Hotsko AL, Robertson S, Mandybur L, Kluger J, White CM. Electrocardiographic and hemodynamic effects of *Panax ginseng*. *Ann Pharmacother*. 2002; 36(5):758–63.
7. Wu W, Zhang XM, Liu PM, Li JM, Wang JF. Effects of *Panax notoginseng* saponin Rg1 on cardiac electrophysiological properties and ventricular fibrillation threshold in dogs. *Zhongguo Yao Li Xue Bao*. 1995; 16(5):459–63.
8. Bai CX, Sunami A, Namiki T, Sawanobori T, Furukawa T. Electrophysiological effects of ginseng and ginsenoside Re in guinea pig ventricular myocytes. *Eur J Pharmacol*. 2003; 476(1):35–44.; [https://doi.org/10.1016/S0014-2999\(03\)02174-5](https://doi.org/10.1016/S0014-2999(03)02174-5)
9. Li M, Ramos LG. Drug-induced QT Prolongation and Torsades de Pointes. *Pharm Ther*. 2017; 42(7):473–7.

201
202
203
204
205
206
207
208
209
210
211

10.

Thwaites BC, Bose M. Very low calorie diets and pre-fast-
ing prolonged QT interval. A hidden potential danger.
West Indian Med J. 1992; 41(4):169–71.

11.

Swenne I, Larsson PT. Heart risk associated with weight
loss in anorexia nervosa and eating disorders: risk fac-
tors for QTc interval prolongation and dispersion. Acta
Paediatr Oslo Nor. 1999; 88(3):304–9.; [https://doi.
doi:10.1080/08035259950170079](https://doi.
doi:10.1080/08035259950170079)

12.

Weinsier RL. Fasting—a review with emphasis on the
electrolytes. Am J Med. 1971; 50(2):233–40.; [https://doi.
doi:10.1016/0002-9343\(71\)90152-5](https://doi.
doi:10.1016/0002-9343(71)90152-5)

13.

Puckett L. Renal and electrolyte complications in eating
disorders: a comprehensive review. J Eat Disord. 2023;
11:26.; <https://doi.doi:10.1186/s40337-023-00751-w>

14.

Jamir SL, Deb CR, Jamir NS. Studies on reproductive
biology and seed biology of panax Pseudoginseng wall.
(Araliaceae): a threatened medicinal plant. Int J Conserv
Sci. 2016;7(4):1127–34. Available form: [https://ijcs.ro/
public/IJCS-16-61_Jamir.pdf](https://ijcs.ro/
public/IJCS-16-61_Jamir.pdf)

212
213
214
215
216
217
218
219

Summary of the case

1	Patient (gender, age)	17, male
2	Final diagnosis	Osteopoikilosis in a patient with orthopedic trauma.
3	Symptoms	Proximal phalanx fracture with degloving of the index finger
4	Medications	Surgery
5	Clinical procedure	Surgery
6	Specialty	Traumatology and orthopedics