




Figure 1. A: First coronary arteriography – RCA, B: RCA after thrombus aspiration, C: RCA after 48 hours on Tirofiban, D: RCA after 1 month on triple anti-thrombotic therapy, E: IVUS Intraluminal thrombus, F: IVUS - Post PCI stent placement, G: RCA after final PCI.

References

1. Vela CNC, Gamarra-Valverde NN, Inga KE, Durand P. Manual thrombus aspiration in patients with ST-elevation myocardial infarction with high thrombus burden and a total ischemic time greater or equal than 3 hours: mini review. *Curr Probl Cardiol*. 2023;48(10):101786. <https://doi.org/10.1016/j.cpcardiol.2023.101786>
2. Pradhan A, Bhandari M, Vishwakarma P, Sethi R. Deferred stenting for heavy thrombus burden during percutaneous coronary intervention for ST-elevation MI. *Eur Cardiol Rev*. 2021;16:e08. <https://doi.org/10.15420/ecr.2020.31>
3. Groenland FTW, Neleman T, Kakar H, Scoccia A, Ziedses des Plantes AC, Clephas PRD, et al. Intravascular ultrasound-guided versus coronary angiography-guided percutaneous coronary intervention in patients with acute myocardial infarction: a systematic review and meta-analysis. *Int J Cardiol*. 2022;353:35–42. <https://doi.org/10.1016/j.ijcard.2022.01.021>