6

10

11

12

13

14 15

16

17

19

20

21 22

23

24

25

26

27

28

29

30

31

32

33

34

35 36

37

38

39

Ovarian teratoma with bowel fistula managed non-operatively

- Thomas Kirengo^{1*}, Samra Ali¹, Samuel Ghattas¹,
- Mostafa AbdelKarim¹, Mohiuddin Kamal¹, Sabina
- Kondokhar², Joseph Mechery², Ali Murtada¹

European Journal of Medical Case Reports

Volume XX(XX):01-06 DOI: XXXX





This is an open access article distributed in accordance with the Creative Commons Attribution (CC BY 4.0) license: https://creativecommons.org/licenses/by/4.0/) which permits any use, Share — copy and redistribute the material in any medium or format, Adapt — remix, transform, and build upon the material for any purpose, as long as the authors and the original source are properly cited. © The Author(s) 2025

ABSTRACT

7 Background: Mature ovarian cystic teratomas (MOCT) are the most common benign ovarian tumours. They are mainly asymptomatic (60%) but can, in rare cases (<1%), complicate by forming a fistula into adjacent structures such as the bowel. 9 Patients with fistulating ovarian teratomas are typically managed surgically, most often requiring laparotomy.

Case Presentation: We report a case of an 18-year-old woman who presented with vague abdominal and gynaecological symptoms. She expelled a mass with hair and teeth per rectum, which was later confirmed as a MOCT. We managed her conservatively in collaboration with the colorectal and gynaecology teams. At one-year follow-up, she remained asymptomatic with no evidence of a residual fistula.

Conclusion: This case highlights that while most MOCTs with bowel fistulation require surgical management, carefully selected patients may be managed conservatively with close follow-up.

Keywords: Mature ovarian cystic teratomas (MOCT), bowel fistula, case report, non-operative treatment.

Type of Article: CASE REPORT Specialty: General Surgery,

Gynaecology

Correspondence to: Thomas Kirengo

*General Surgery Department, Ysbyty Glan Clwyd, Rhyl, UK.

Email: Thomas.kirengo@nhs.wales.uk

Full list of author information is available at the end of the article.

Received: 12 August 2025 Revised (1): 24 September 2025

Accepted: 30 September 2025

Background

Ovarian teratoma, a germ cell tumour, is the most common benign ovarian tumour [1]. Mature teratomas, synonymously referred to as mature ovarian cystic teratoma (MOCT) or dermoid cysts, account for >95% of all teratomas and 20%-25% of all ovarian tumours [2, 3]. MOCTs contain well-differentiated tissues from all three germ cell layers (ectoderm, mesoderm, and endoderm), macroscopically seen with structures such as teeth, hair, and bones

MOCTs are typically asymptomatic in approximately 60% of cases, and are often discovered incidentally on imaging for unrelated clinical indications. However, symptomatic patients can complain of abdominopelvic pain or masses. Given the vague symptoms, these patients may present to either gynaecologists or general surgeons [2]. This is further complicated when patients develop a fistula and present with unusual clinical symptoms such as passing hair in stool [5].

Fistula formation in MOCTs is a rare complication seen in fewer than 1% of all cases [3]. The underlying pathophysiology is, however, not well studied [6]. In a series of 17 cases, Kizaki et al. reported of fistula formation involving MOCTs and adjacent organs, such as the bladder and bowel. They identified inflammation (76.5%) as the most common cause, followed by malignant transformation (23.5%). All these cases were managed with laparotomy [3].

42

43

44

45

46

47

48

49

50

51

52

54

55

56

57

59

60

Patients with fistulating ovarian teratomas generally require laparotomy for surgical repair. Here, we report a case of an 18-year-old woman who spontaneously passed a MOCT per rectum and was managed conservatively. To the best of our knowledge, this represents the only documented instance of successful non-operative management to date.

Case Presentation

An 18-year-old nulliparous woman presented initially with intermittent right iliac fossa pain and copious yellow vaginal discharge. She had an ultrasound scan done at that time, which revealed the presence of a large lesion of unknown aetiology on the midline and right adnexa. In addition, it was noted that her copper IUCD was incorrectly positioned just above the cervix. She was reviewed by the Gynaecologists, the Copper IUCD was removed, a vaginal swab was taken for microscopy, culture, and sensitivity (no organisms seen), and follow-up review after MRI scan of the abdomen and pelvis was arranged.

However, three months later, the patient re-presented to the emergency department with left iliac fossa pain, per rectal bleeding, and diarrhoea. Her pain was not associated with nausea, vomiting, or fever. Physical examinations revealed no signs of peritonitis. Blood test showed leucocytosis with (WBC 12.2 x10 9 /L; reference 4.0-11.0), thrombocytosis (platelet count 658 x10 9 /L; reference 150-400), neutrophilia (8.5 x10 9 /L; reference 1.7-7.5), prolonged PT (13.0 sec; reference 9.0-12.0), elevated CRP (80 mg/L; reference <5) and raised CA 125 (75 KU/L; reference <35). Other tumour markers were within normal ranges, notably β -HCG (<1 IU/L; reference <1), Alphafetoprotein (2 KU/L; reference 1-6), and CEA (1 ug/L; reference <4).

Urgent Imaging was requested; however, on the same day, she spontaneously expelled a mass per rectum. Gross morphological inspection demonstrated well-developed tissues, including teeth and hair (Figure 1).

Investigations

CT scan done on her second presentation revealed a sizable pelvic collection with evidence of a large wall defect involving the distal sigmoid/rectosigmoid junction with surrounding active inflammatory change and free fluid (Figure 2). Retrospective review of the scans at the Gynaecology MDT, a month later, identified evidence



Figure 1. Image of gross expelled mass showing teeth and hair

of material within the bowel (teeth) compatible with a dermoid.

Furthermore, an MRI pelvis was also conducted three days after the CT-scan, which showed a moderate volume of partially loculated pelvic free fluid and a 6.4 x 5.9 cm heterogeneous, gas-containing central pelvic collection communicating with a wide fistulous defect within the anterior wall of the mid sigmoid colon, which appeared to be left Tubo-ovarian in origin (Figure 3).

Histopathology analysis two weeks later confirmed the presence of teeth, hair, sebaceous glands, and concluded a diagnosis of mature ovarian teratoma.

Differential Diagnosis

Our differential diagnoses evolved as the patient's symptoms changed. At the first presentation, pelvic inflammatory disease with a tubo-ovarian abscess was considered, given the vaginal discharge and ultrasound findings of a malpositioned intrauterine device. Following the development of left iliac fossa pain and rectal bleeding, colorectal pathology such as colitis or inflammatory bowel disease was entertained. Ultimately, the passage of tissue per rectum, together with cross-sectional imaging and histology, confirmed the diagnosis of a fistulating ovarian teratoma.

Treatment

The patient was not keen on any invasive treatment or investigations, which guided our choice of management. She was managed conservatively in collaboration with the Colorectal team and discharged following a 5-day inpatient stay. She remained asymptomatic and returned back to regular activity within two months.

Outcome and Follow-up


Follow-up MRI scans at one year demonstrated no definitive evidence of a residual fistula. Although she initially declined endoscopy, she subsequently consented to flexible sigmoidoscopy for further evaluation, which at 21 months identified scarring at the site of the previous sigmoid fistula with no evident fistula opening (Figure 4). The patient was reassured.

Discussion

Ovarian teratoma with bowel fistulation is a rare condition, described in both paediatric and elderly patients, with the youngest documented case reported in a 9-year-old girl and the oldest in an 85-year-old woman [7, 8]. The first reported case in literature was in 1965 [9]. Our patient was an otherwise healthy young woman, under 20 years of age, who presented with vague abdominal and gynaecological symptoms. Importantly, this case is distinctive in documenting successful conservative management, with no apparent complications observed at 21 months follow-up.



Figure 2. CT-scan – Dermoid within the sigmoid colon. Sagittal View: Small Blue Arrow demonstrates Dermoid in Rectum; Large Blue Arrow shows fistula in sigmoid colon. Coronal View: Large Blue arrow highlights the Dermoid cyst.



142

145

146

147

148

149

150

151

152

153

154

155

156

Figure 3. MRI T2 Sagittal sequence- Dermoid passed, but fistula into the sigmoid evident as shown by the blue arrow.

Approximately 80% of females diagnosed with ovarian teratomas are of reproductive age [10]. Abdominal pain is one of the most frequently reported presenting symptoms [11]. Other manifestations reported include gastrointestinal symptoms such as rectal bleeding, passage of hair with stool, diarrhoea, and tenesmus [12]. Genitourinary symptoms may be found in cases involving the urinary system. These include urinary frequency, dysuria, and haematuria [5]. Unfortunately, the vague and heterogeneous nature of these symptoms makes diagnosis challenging. Our patient

initially presented with right iliac fossa pain, which later localised on the left side, underscoring the potential for misdiagnosis, particularly if confounding factors and red herrings such as a misplaced IUCD are present. Radiological imaging and histology are essential in confirming the diagnosis.

157

158

159

160

161

162

163

164

165

166

167

169

170

171

172

173

175

176

177

178

179

180

181

182

183

184

185

Choice of imaging depends on local availability. Ultrasound is usually considered first line due to ease of access and minimal invasiveness [13]. CT scan has higher sensitivity (93%-98%) compared to ultrasound, and provides detailed cross-sectional images, enabling identification of characteristic features and relationship to surrounding viscera [13, 14]. Additionally, MRI offers superior soft tissue resolution, detecting features potentially missed on CT [13]. In our case report, an emergency CT revealed a distal sigmoid colon wall fistula defect and pelvic collection. This was subsequently confirmed on the MRI of the pelvis. Although initially missed, the fistulated teratoma was retrospectively seen on review of the CT scan at an MDT meeting. Endoscopy, although invasive, enables direct visualisation and tissue diagnosis. Flexible sigmoidoscopy was later performed despite initial resistance from our patient. Overall, multimodal imaging is necessary to enhance diagnostic accuracy and identify complications.

Ovarian teratomas that fistulate into the bowel present a complex clinical scenario. The majority of published case reports involve surgical treatment with laparotomy or laparoscopy. This is possibly due to the extent of




Figure 4. Distal Sigmoid Flexible sigmoidoscopy.

pathology, the need for direct visualisation, and histological assessment [3]. To the best of our knowledge, this case report is the only documented instance of conservative management for fistulating disease. Standard surgical management involves excision of the teratoma, repair or resection of affected bowel or adjacent structures, and commonly unilateral or bilateral salpingo-oopherectomy. Therefore, young patients should be counselled on the risk of loss of fertility. Our patient was 18 years old at the time of presentation and was not keen on invasive treatment. Additionally, our patient fully expelled the dermoid in whole, as later confirmed by MRI imaging. Conservative treatment, however, carries risks such as the malignant transformation into squamous cell carcinoma, which is associated with poor prognosis [8, 10]. For this reason, close follow-up with imaging was instituted to detect any complications early. A multidisciplinary approach involving surgeons, radiologists, gynaecologists, and other relevant specialists is essential to optimise clinical outcomes.

Limitations

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

211

212

213

214 215

216

217

218

219

220

221

Patient follow-up was limited to 21 months. Management 208 was influenced by patient preference, and the rarity of the 209 210 condition limits comparative analysis.

Conclusion

This case highlights that while most MOCTs with bowel fistulation require surgical management, carefully selected patients may be managed conservatively with close follow-up. This points to the importance of individualised multidisciplinary care, taking into consideration patient factors such as age and patient wants. A rigorous systematic review is recommended to identify a wider range of cases and management options.

Although most MOCTs with bowel fistulation require surgery, this case demonstrates that carefully selected

patients may be managed conservatively with close surveillance. Individualised multidisciplinary care, considering patient age and preferences, is essential. A systematic review is warranted to provide a comprehensive evaluation of the available literature on this condition.

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

248

249

250

251

What Is New?

- Ovarian teratoma with bowel fistulation is a rare complication, reported in women ranging in age from as young as 9 years to 85 years old.
- Clinical presentation can be vague, with overlapping gastrointestinal and gynaecological symptoms, which might delay timely diagnosis.
- Passage of tissue containing hair or teeth per rectum is pathognomonic of a mature ovarian cystic teratoma and should raise the index of suspicion for fistulating disease.
- · Cross-sectional imaging and histopathology are essential for confirming the diagnosis and defining the extent of fistulation
- Cases reported in the literature have been managed surgically; to the best of our knowledge, this is the first case demonstrating conservative management.

List of Ahhreviations

ist of Abbreviations:		
CEA	Carcinoembryonic Antigen	
HCG	Human Chorionic Gonadotropin	245
MOCT	Mature Ovarian Cystic Teratoma	246
PT	Prothrombin Time	247

Conflict of interests

The authors declare that there is no conflict of interest regarding the publication of this article.

Funding

None	252

253 Consent for publication

- 254 Informed written consent was given by the patient for the publi-
- 255 cation of this case report and accompanying images. All patient
- 256 identifiers were removed to ensure anonymity.

257 Ethical approval

- 258 The authors assessed the case in line with the UK National
- 259 Research Ethics requirements. Formal Ethical approval is not
- 260 required at our institution to publish an anonymous case report.

261 Author details

- 262 Thomas Kirengo¹, Samra Ali¹, Samuel Ghattas¹, Mostafa
- 263 AbdelKarim¹, Mohiuddin Kamal¹, Sabina Kondokhar², Joseph
- 264 Mechery², Ali Murtada¹
- 265 1 General Surgery Department, Ysbyty Glan Clwyd, Rhyl, UK
- 266 2 Gynaecology Department, Ysbyty Glan Clwyd, Bodelwyddan,267 Rhyl, UK

268 References

- 269 1. Haque R, Akter T. Clinical presentation and histological 270 types of ovarian tumor in patients admitted in tertiary 271 care center. Sch Int J Obstet Gynec. 2023;6(11):450–6. 272 https://doi.org/10.36348/sijog.2023.v06i11.004
- Cong L, Wang S, Yeung SY, Lee JHS, Chung JPW, Chan DYL.
 Mature cystic teratoma: an integrated review. Int J Mol Sci.
 2023;24(7):141. https://doi.org/10.3390/ijms24076141
- Kizaki Y, Nagai T, Ohara K, Gomi Y, Akahori T, Ono Y, et al.
 Ovarian mature cystic teratoma with fistula formation into the rectum: a case report. Springerplus. 2016;5(1):1700. https://doi.org/10.1186/s40064-016-3426-4
- 280 4. Chen VW, Ruiz B, Killeen JL, Coté TR, Wu XC, Correa CN.
 281 Pathology and classification of ovarian tumors. Cancer.
 282 2003;97(10 Suppl):2631–42. https://doi.org/10.1002/
 283 cncr.11345
- Singh R, Joshi S, Hatimi H, Somani P, Rath P, Joshi RM. Rectal fistula due to ovarian teratoma. Endoscopy. 2012;44 Suppl
 UCTN:E260. https://doi.org/10.1055/s-0032-1309394

 Conway AZ, Hanna T. A dermoid cyst fistulating with the transverse colon. Ann R Coll Surg Engl. 2012;94(7):e230–1. https://doi.org/10.1308/003588412X13373405385935 287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

- Dandia SD. Rectovesical fistula following an ovarian dermoid with recurrent vesical calculus: a case report.
 J Urol. 1967;97(1):85–7. https://doi.org/10.1016/ S0022-5347(17)62984-7
- Chong HM, Lee FY, Lo A, Li CM. A giant gas-filled abdominal mass in an elderly female: a case report. World J Gastroenterol. 2011;17(31):3659–62. https://doi.org/10.3748/wig.v17.i31.3659
- Figiel LS, Figiel SJ. Report of two cases of sigmoidal fistula complicating tubo-ovarian disease. Dis Colon Rectum. 1966;9(2):107–8. https://doi.org/10.1007/BF02617308
- Agrawal N, Gothwal M, Yadav G, Singh P, Varshney V, Yadav T, et al. Ileo-dermoid fistula: a rare presentation of ovarian dermoid. J Midlife Health. 2023;14(4):299–301. https://doi.org/10.4103/jmh.jmh_205_22
- 11. Farkouh E, Allard M, Paquin JG. Benign solid teratoma of the ovary with rupture into the rectum. Can J Surg. 1982;25(1):77–8.
- Rakower SR, Schinella RA, Fazzini EP. Benign solid teratoma of the fallopian tube with rupture into the rectum: report of a unique rectal tumor. Dis Colon Rectum. 1976;19(2):167–71. https://doi.org/10.1007/BF02590875
- Saleh M, Bhosale P, Menias CO, Ramalingam P, Jensen C, Iyer R, et al. Ovarian teratomas: clinical features, imaging findings and management. Abdom Radiol. 2021;46(6):2293–307. https://doi.org/10.1007/s00261-020-02873-0
- 14. Saba L, Guerriero S, Sulcis R, Virgilio B, Melis G, Mallarini G. Mature and immature ovarian teratomas: CT, US and MR imaging characteristics. Eur J Radiol. 2009;72(3):454–63. https://doi.org/10.1016/j.ejrad.2008.07.044

Summary of the case

322

1	Patient (age, gender)	18 years, female				
2	Final diagnosis	Mature ovarian teratoma with bowel fistula				
3	Symptoms	Right iliac fossa pain, yellow vaginal discharge, later left iliac fossa pain, per rectal bleeding, diarrhoea				
4	Medications	N/A				
5	Clinical Procedure	Conservative management – no surgery performed				
6	Specialty	General Surgery, Gynaecology				