

Tonsillar trichophagia without trichotillomania: a pediatric case and psychiatric considerations

Abdullah Ozan Polat^{1*} , Okcan Basat¹ 

European Journal of Medical Case Reports

Volume 10(1):35–36

DOI: 10.24911/ejmcr.9-2391



This is an open access article distributed in accordance with the Creative Commons Attribution (CC BY 4.0) license: <https://creativecommons.org/licenses/by/4.0/> which permits any use, Share — copy and redistribute the material in any medium or format, Adapt — remix, transform, and build upon the material for any purpose, as long as the authors and the original source are properly cited. © The Author(s) 2026

Type of Article: LETTER TO THE EDITOR **Specialty:** Psychiatry

Correspondence to: Abdullah Ozan Polat

*Family Physician Specialist, Department of Family Medicine, University of Health Sciences, Gaziosmanpaşa Training and Research Hospital, Istanbul, Türkiye. Istanbul/Turkey.

Email: ozanpolat1987@gmail.com

Full list of author information is available at the end of the article.

Received: 15 October 2025

Revised (1): 11 November 2025

Accepted: 13 November 2025

Letter to Editor

Dear Editor,

Trichophagia, defined as the compulsive ingestion of hair, is a rare but clinically significant condition most often associated with trichotillomania [1]. However, isolated trichophagia without accompanying hair-pulling behavior is exceedingly uncommon, particularly in pediatric populations [2]. We report a rare case of a nine-year-old girl diagnosed with isolated tonsillar trichophagia, emphasizing its psychiatric implications and the importance of multidisciplinary management.

A nine-year-old girl presented with a one-week history of sore throat and dysphagia. Physical examination revealed dark, brush-like hair deposits embedded within

the bilateral tonsillar crypts (Figure 1A). There was no alopecia or self-reported hair-pulling behavior. Her mother had obsessive-compulsive disorder, and her father had generalized anxiety disorder, suggesting strong familial psychiatric vulnerability [3]. Tonsillectomy confirmed hair bezoars localized within the tonsillar crypts (Figure 1B). Postoperatively, the patient was referred to a child and adolescent psychiatry clinic for cognitive-behavioral therapy [4] and family-based psychoeducation.

This case underscores that trichophagia can exist independently of trichotillomania and may manifest in unusual anatomical sites such as the tonsils [5]. Clinicians should maintain high diagnostic awareness and consider psychiatric evaluation even when trichophagia occurs without




Figure 1. (A) Dark, brush-like hair deposits embedded within the bilateral tonsillar crypts. (B) Post-tonsillectomy view showing complete removal

overt hair-pulling. Early psychiatric intervention, combined with surgical management, remains essential to prevent recurrence and address underlying psychopathology.

Acknowledgments

None.

Conflict of Interest

The authors declare no conflicts of interest.

Funding

None.

Consent to participate

The study was conducted according to the guidelines of the Declaration of Helsinki. Written informed consent was obtained from the participant included in the study.

Ethical approval

Ethical approval is not required for anonymous case reports at our institution. Our institution (Gaziosmanpaşa Training and Research Hospital) does not require ethical approval for reporting individual cases or case series. Our study is in accordance with the Declaration of Helsinki and CARE guidelines. Written informed consent for publication of identifying images or other personal or clinical details was obtained from the patient. The photographs are completely unidentified, and personal details are not mentioned in the text. The authors are accountable for all aspects of the work and for ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Informed consent

Written informed consent was obtained from the patient's parents for publication of this case and accompanying images.

Data availability statement

All data generated or analysed during this study are included in this published article. The study data were stored. The data used and analyzed during the current study could be available from the corresponding author upon reasonable request.

Author contributions

All of the authors declare that they have all participated in the design, execution, and analysis of the paper, and that they have approved the final version.

Author details

Abdullah Ozan Polat¹, Okcan Basat¹

1. Health Sciences University Gaziosmanpaşa Training and Research Hospital, Family Medicine Clinic, Istanbul, Turkey

References

1. Grant JE, Chamberlain SR. Trichotillomania and skin-picking disorder: an update. *Focus. Am Psychiatr Publ.* 2021;19(4):405–12. <https://doi.org/10.1176/appi.focus.20210013>
2. Schuler L, Hodel M, Stieger C. The Rapunzel syndrome: a hairy tale. *Surg Case Rep.* 2023;9(1):49. <https://doi.org/10.1186/s40792-023-01631-w>
3. Chacon P, Bernardes E, Faggian L, Batistuzzo M, Moriyama T, Miguel EC, et al. Obsessive-compulsive symptoms in children with first degree relatives diagnosed with obsessive-compulsive disorder. *Braz J Psychiatry.* 2018;40(4):388–93.
4. Bortoncello CF, Cardoso NO, Xavier SRM, Ferrão YA. Effectiveness of online group cognitive behavioral therapy for adolescents with obsessive-compulsive disorder: a pilot study. *Braz J Psychiatry.* 2024;46:e20233376. <https://doi.org/10.47626/1516-4446-2023-3376>
5. Flagstad E, Hummelen B, Moen E, Dammen T, Haaland Å, Sunde T, et al. Early maladaptive schemas in trichotillomania and skin-picking disorder: their relationships with symptom severity and subtypes. *BMC Psychol.* 2025;13(1):789. <https://doi.org/10.1186/s40359-025-03096-y>