

68 AICN occurs due to the obstruction of the ureter and/
69 or renal tubules by AC [7]. It is thought to be related to
70 a concentration-dependent oversaturation of amoxicil-
71 lin and subsequent crystallization in the urine following
72 glomerular filtration and tubular secretion of amoxicillin
73 [1,10]. Risk factors for the crystallization and tubular de-
74 position of amoxicillin are related to the pathophysiology
75 and include administration of high doses (> 12 g/d), rapid

76 infusion (> 2 g/30 min), oliguria, dehydration, hypo-
77 volemia, and low urinary pH [10,11].

78 Treatment consists of discontinuation of amoxicillin
79 treatment and improving tubular flow and urine output
80 with volume resuscitation [2]. About 70%-80% of the
81 patients developing AC experience renal failure [11].
82 However, the prognosis is mostly favorable, and patients
83 generally recover to normal kidney function 3-17 days




Figure 1. Schematic overview of case.

84 after stopping amoxicillin [1,2,10]. Still, temporary renal
85 replacement therapy is needed in 10%-40% of patients [2].

86 Case Description

87 A 75-year-old woman with no relevant medical history
88 presented herself to the emergency department with high
89 fever and general malaise (Figure 1). She reported sev-
90 eral days of poor appetite and a dry mouth, with complete
91 absence of oral intake. In particular, she did not recog-
92 nize that she had a fever. She denied respiratory, gastro-
93 intestinal, or urinary symptoms. Her left leg, which had
94 long-standing varicose veins, showed new-onset redness
95 and was painful to touch. On clinical examination, her
96 vital signs were notable for a fever of 40.2°C. She was
97 hemodynamically stable. Cardiac exam revealed no mur-
98 murs, but she was tachycardic with jugular venous dis-
99 tension. Lung auscultation and abdominal examination
100 were normal. Lower extremities exhibited bilateral pitting
101 edema, more pronounced on the left, with erythema and
102 tenderness. Neurological examination was unremarkable.

103 Initial work-up and bedside ultrasound suggested
104 extensive cellulitis of the left lower leg without evidence
105 of deep vein thrombosis. Other initial investigations (e.g.,
106 X-ray of chest, abdominal ultrasound) were negative, so
107 at that time the cellulitis seemed to be the only identifiable
108 focus of fever. Empiric intravenous amoxicillin/clavulanic
109 acid was initiated in combination with IV hypotonic fluids
110 because of the absence of oral intake. Blood cultures later
111 yielded *Streptococcus dysgalactiae*, which was also iso-
112 lated from a wound culture from the affected leg. Due to
113 persistent fever, a transesophageal echocardiogram (TEE)
114 was performed, which confirmed native aortic valve endo-
115 carditis with vegetations on the non-coronary cusp. In line
116 with the 2023 European Society of Cardiology guidelines,
117 antibiotic therapy was adjusted to high-dose IV amoxi-
118 cillin (6 × 2 g/day) [12]. In addition, the TEE confirmed
119 intravascular underfilling, for which the ongoing hypo-
120 tonic intravenous fluid infusion rate was increased.

121 Two days later, the patient developed progressive AKI,
122 with macroscopic hematuria and non-nephrotic range pro-
123 teinuria. Urinary pH was in the low-normal range. Initial
124 differential diagnosis included prerenal kidney injury given
125 the intravascular underfilling on TEE, acute tubular necro-
126 sis possibly linked to a short hypotensive episode, sepsis-in-
127 duced AKI, acute interstitial nephritis (AIN) secondary
128 to antibiotic exposure, septic emboli, and peri-infectious
129 glomerulonephritis. Despite supportive care - including
130 adjustment of the intravenous fluid infusion rate and admin-
131 istration of sodium bicarbonate - the renal function deterio-
132 rated further, necessitating acute hemodialysis.

133 The differential diagnosis was approached as followed:
134 prerenal kidney injury became less likely after correct fluid
135 resuscitation failed to improve kidney function. Sepsis-
136 induced AKI was not favored, as macroscopic hematuria is
137 uncommon in this condition. AIN was deemed less likely,




Figure 2. Digital light microscopy (Atellica UAS8000 of Siemens Healthineers) revealing amoxicillin crystals: needle-shaped elongated crystals A: Digital light microscopy (Atellica UAS800 of Siemens Healthineers) revealing AC: a needle-shaped elongated crystal. B: Digital light microscopy (Atellica UAS800 of Siemens Healthineers) revealing AC: two needle-shaped elongated crystals and large aggregated needle-shaped crystals. C: Digital light microscopy (Atellica UAS800 of Siemens Healthineers) revealing AC: two needle-shaped elongated crystals.

given the early onset of AKI after amoxicillin exposure. 138
Embolic phenomena were unlikely, since clinically signifi- 139
cant renal dysfunction would require bilateral involvement, 140

141 which is rather uncommon. Eventually, the main diagnostic
 142 challenge remained distinguishing peri-infectious glomeru-
 143 lonephritis from possible crystal nephropathy.

144 Given this uncertainty and the further deterioration of
 145 kidney function, urine microscopy was repeated 4 days
 146 after the onset of AKI, revealing the presence of elongated
 147 crystals, which confirmed the diagnosis of AICN (Figure
 148 2A-C). Amoxicillin was discontinued and changed to
 149 Penicillin G in continuous infusion. After discontinua-
 150 tion of amoxicillin in favor of intravenous penicillin, the
 151 kidney function improved rapidly within the next 7 days
 152 and eventually returned to baseline (Figure 3). This rapid
 153 improvement of renal function also ruled out acute tubu-
 154 lar necrosis as the cause of AKI. Additional renal biopsy
 155 was considered obsolete, given its invasiveness, bleeding
 156 risk, and the prior confirmation of typical AC along with
 157 rapid improvement after drug withdrawal. Consequently,
 158 temporary hemodialysis was fully withdrawn after several
 159 days (Figure 4 – timeline).

160 Follow-up echocardiography at 2 weeks showed stable
 161 non-progressive aortic insufficiency, with persistent
 162 vegetations on the aortic valve, indication for a further
 163 conservative approach. After 4 weeks of intravenous anti-
 164 biotic treatment, nevertheless, she developed progressive

165 aortic regurgitation necessitating aortic valve replace-
 166 ment. Renal function remained stable during this period,
 167 and monthly follow-up over 1 year after discharge con-
 168 firmed persistently normal values.

169 **Discussion**

170 AICN is an under recognized, but important complication
 171 of a commonly prescribed antibiotic. Our patient had sev-
 172 eral recognized risk factors for AICN, including high-dose
 173 amoxicillin, dehydration, and poor oral intake. Preventive
 174 measures such as the administration of intravenous hypo-
 175 tonic maintenance fluids and adjusting the infusion rate in
 176 response to intravascular underfilling on TEE were imple-
 177 mented before the onset of AKI, while urine alkalinization
 178 was initiated after AKI-onset. In retrospect, optimization
 179 of intravascular volume in the context of hypotension and
 180 persistent intravascular underfilling might have been bet-
 181 ter achieved by adding isotonic resuscitation fluids rather
 182 than simply increasing the infusion rate of hypotonic
 183 maintenance fluids. This better management of hypo-
 184 volemia could potentially have reduced the severity of
 185 acute renal injury.

186 Although historically considered rare, different studies
 187 now suggest an increase in the incidence of AICN [3,6,7].




Figure 3. Serum creatinine as a function of time.




Figure 4. Timetable summarizing antibiotic exposure, onset of AKI, timing of urine microscopy, dialysis and recovery.

188 The potential impact of reporting bias and increased aware-
 189 ness on this rise in reports should, however, be acknowl-
 190 edged, given the fact that the most recent data come from
 191 pharmacovigilance studies conducted in France. Thus,
 192 its true incidence remains unclear, also considering the
 193 absence of standardized diagnostic criteria and underuse
 194 of crystal identification through urine microscopy [3,6,7].
 195 The pharmacovigilance studies propose three criteria for
 196 the diagnosis of AICN - AKI, temporal association with
 197 amoxicillin use, and micro- or macroscopic hematuria
 198 [3,6,7]. However, these criteria have limitations: hema-
 199 turia is not consistently present, and the concept of ‘tem-
 200 poral association’ is poorly defined, as AKI may develop
 201 immediately or several days after initiation of treatment.
 202 Moreover, it makes it difficult to differentiate from infec-
 203 tious glomerulonephritis, which would also present with
 204 (albeit glomerular) hematuria and AKI in temporal associ-
 205 ation with infection (and thus antibiotic use).

206 Another complicating factor in diagnosing AICN is the
 207 differential diagnosis with sepsis-induced AKI and AIN.
 208 Sepsis-induced AKI often poses the greatest diagnostic
 209 challenge in infectious contexts due to the lack of specific
 210 biomarkers distinguishing it from other forms of AKI [2].
 211 It is typically accompanied by dysfunction of at least one
 212 other organ system and usually develops within 7 days of
 213 sepsis onset [2]. In contrast, AIN represents a hypersensi-
 214 tivity reaction frequently triggered by drugs such as amox-
 215 icillin, generally occurring after a longer latency (>10
 216 days) and may rarely be accompanied by fever, rash, and
 217 arthralgia [9]. It should be suspected in patients with ris-
 218 ing serum creatinine and urinalysis showing white blood
 219 cells, white cell casts, or eosinophiluria, although urinary
 220 findings may be minimal [13]. Recovery from AIN is typi-
 221 cally slower than from AICN, and a definitive diagnosis of
 222 either condition requires a kidney biopsy [2,9].

223 In clinical practice, biopsy is rarely performed if the
 224 diagnosis for underlying AKI is strongly supported by
 225 clinical context, if renal function improves after stopping
 226 the suspected drug, and if urine microscopy provides suf-
 227 ficient evidence when typical needle-shaped crystals are
 228 identified. All three conditions were present in this case,
 229 supporting a presumptive but well-supported diagnosis of
 230 AICN [7].

231 Although urine microscopy is widely available, reli-
 232 able detection of AC requires prompt examination of fresh
 233 urine with polarized light microscopy and operator exper-
 234 tise, as the characteristic needle- or rod-like crystals can
 235 be easily missed. False negatives are common, particu-
 236 larly with delayed processing, transient crystalluria, anu-
 237 ria, crystal dissolution in alkaline urine, or milder cases.
 238 Nonetheless, urine microscopy remains a rapid diagnostic
 239 tool, especially when the lack of adequate diagnosis could
 240 cause a delay in appropriate therapy [14].

241 This case report is relevant as it highlights a relatively
 242 rare but important complication of a commonly used drug.
 243 It illustrates that even with limited diagnostics, the diagno-
 244 sis of AICN can be confidently made, allowing for prompt
 245 intervention and rapid reversal of AKI. A key component
 246 in this process is urine microscopy, which can detect AC
 247 and aid in differentiating AICN from other conditions, as
 248 demonstrated in this case report. Nonetheless, as a sin-
 249 gle-case report, its findings remain inherently limited and
 250 cannot be generalized.

Conclusion

251 AICN is an unusual but potentially severe complication of
 252 high-dose intravenous amoxicillin. This case demonstrates
 253 that in the presence of a broad differential diagnosis, the detec-
 254 tion of typical urinary crystals through urine microscopy,
 255 together with the temporal relationship between amoxicillin
 256 use and AKI, can provide sufficient diagnostic confidence,
 257 without the need for invasive testing. Confirmation of AC
 258 via urine microscopy was the key element in guiding timely
 259 therapeutic decisions. Early recognition facilitated by urine
 260 microscopy is crucial, as discontinuation of amoxicillin usu-
 261 ally leads to rapid and complete renal recovery.
 262

What is new?

263 This case emphasizes the importance of urine microscopy as
 264 a rapid, non-invasive, and accessible diagnostic tool for AICN,
 265 providing clear evidence that may avoid the need for kidney
 266 biopsy. The detection of characteristic amoxicillin crystals
 267 allows clinicians to narrow a broad differential diagnosis and
 268 guide immediate, appropriate therapeutic interventions,
 269 supporting the value of its routine use in the evaluation of
 270 acute kidney injury during amoxicillin treatment.
 271

272 **List of Abbreviations**

273	AC	Amoxicillin-crystals
274	AICN	Amoxicillin-induced crystal nephropathy
275	AIN	Acute interstitial nephritis
276	AKI	Acute kidney injury
277	TEE	Transesophageal echocardiogram

278 **Conflict of interest**

279 *The authors declare that they have no conflict of interest regard-*
 280 *ing the publication of this case report.*

281 **Funding**

282 None.

283 **Consent for publication**

284 The patient provided written informed consent for the publica-
 285 tion of the clinical details and any associated images included in
 286 this case report.

287 **Ethical approval**

288 The Ethics Committee of the General Hospital AZ Sint-Lucas
 289 Brugge raised no objection to publication. Ethical approval date:
 290 06-10-2025; letter number AZSTL2025-013.

291 **Author details**

- 292 Lore Bogaert¹, Roos Pannecoek², Lisa Florin³, Dirk Verleyen⁴
- 293 1. Internal Medicine, UZ Leuven, Bruges, Belgium
- 294 2. Nephrology, AZ Sint-Lucas, Bruges, Belgium
- 295 3. Laboratory Medicine, AZ Sint-Lucas, Bruges, Belgium
- 296 4. Cardiology, AZ Sint-Lucas, Bruges, Belgium

297 **References**

298 1. Couto J, Pontes Dos Santos L, Carlos Alves J, López
 299 R, Maldonado C. Amoxicillin crystalluria: a rare side
 300 effect of a commonly prescribed antibiotic. *Eur J*
 301 *Case Rep Intern Med*. 2017;4(10):736. [https://doi.](https://doi.org/10.12890/2017_000736)
 302 [org/10.12890/2017_000736](https://doi.org/10.12890/2017_000736)

303 2. Vodovar D, Mousseaux C, Daudon M, Jamme M,
 304 Letavernier E. Amoxicillin crystalluria and amoxicillin-in-
 305 duced crystal nephropathy: a narrative review. *Kidney*
 306 *Int*. 2025;107(1):33–43. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.kint.2024.09.019)
 307 [kint.2024.09.019](https://doi.org/10.1016/j.kint.2024.09.019)

308 3. Vodovar D, Thomas L, Mongardon N, Lepeule R, Lebrun-
 309 Vignes B, Biour M, et al. Dramatic increase of amox-
 310 icillin-induced crystal nephropathy found in a cohort

study of French Pharmacovigilance Centers. *Antimicrob*
 311 *Agents Chemother*. 2018;62(3):e01630–17. [https://doi.](https://doi.org/10.1128/AAC.01630-17)
 312 [org/10.1128/AAC.01630-17](https://doi.org/10.1128/AAC.01630-17)

313 4. Fogazzi GB. Amoxycillin, a rare but possible cause of
 314 crystalluria. *Nephrol Dial Transplant*. 2003;18(1):212–4.
 315 <https://doi.org/10.1093/ndt/18.1.212>

316 5. Kellum JA, Lameire N. Diagnosis, evaluation, and man-
 317 agement of acute kidney injury: a KDIGO summary (Part
 318 1). *Crit Care*. 2013;17(1):204. [https://doi.org/10.1186/](https://doi.org/10.1186/cc11454)
 319 [cc11454](https://doi.org/10.1186/cc11454)

320 6. Garnier AS, Drablier G, Briet M, Augusto JF. Nephrotoxicity
 321 of amoxicillin and third-generation cephalosporins: an
 322 updated review. *Drug Saf*. 2023;46(8):715–24. [https://](https://doi.org/10.1007/s40264-023-01316-1)
 323 doi.org/10.1007/s40264-023-01316-1

324 7. Thomas L, Le Beller C, Trenque T, Michot J, Zenut M,
 325 Letavernier E, et al. Amoxicillin-induced crystal nephrop-
 326 athy: a nationwide French pharmacovigilance databases
 327 study. *Br J Clin Pharmacol*. 2020;86(11):2256–65. [https://](https://doi.org/10.1111/bcp.14328)
 328 doi.org/10.1111/bcp.14328

329 8. Zeller V, Puyraimond-Zemmour D, Sené T, Lidove O,
 330 Meyssonier V, Ziza JM. Amoxicillin crystalluria, an
 331 emerging complication with an old and well-known anti-
 332 biotic. *Antimicrob Agents Chemother*. 2016;60(5):3248.
 333 <https://doi.org/10.1128/AAC.00359-16>

334 9. Jamme M, Oliver L, Ternacle J, Lepeule R, Moussafer
 335 A, Haymann JP, et al. Amoxicillin crystalluria is associ-
 336 ated with acute kidney injury in patients treated for
 337 acute infective endocarditis. *Nephrol Dial Transplant*.
 338 2021;36(10):1955–68. [https://doi.org/10.1093/ndt/](https://doi.org/10.1093/ndt/gfab074)
 339 [gfab074](https://doi.org/10.1093/ndt/gfab074)

340 10. Mousseaux C, Rafat C, Letavernier E, Frochet V, Kerroumi
 341 Y, Zeller V, et al. Acute kidney injury after high doses of
 342 amoxicillin. *Kidney Int Rep*. 2020;6(3):830–4. [https://doi.](https://doi.org/10.1016/j.ekir.2020.11.040)
 343 [org/10.1016/j.ekir.2020.11.040](https://doi.org/10.1016/j.ekir.2020.11.040)

344 11. Cozzo D, Forni Ognà V, Ventresca M. Amoxicillin
 345 crystalluria during endocarditis treatment. *Kidney*
 346 *Int*. 2025;107(1):194. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.kint.2024.07.029)
 347 [kint.2024.07.029](https://doi.org/10.1016/j.kint.2024.07.029)

348 12. Delgado V, Ajmone Marsan N, De Waha S, Bonaros N, Brida
 349 M, Burri H, et al. 2023 ESC Guidelines for the management
 350 of endocarditis. *Eur Heart J*. 2023;44(39):3948–4042.

351 13. Nussbaum EZ, Perazella MA. Diagnosing acute intersti-
 352 tial nephritis: considerations for clinicians. *Clin Kidney J*.
 353 2019;12(6):808–13. <https://doi.org/10.1093/ckj/sfz080>

354 14. Becker GJ, Garigali G, Fogazzi GB. Advances in urine
 355 microscopy. *Am J Kidney Dis*. 2016;67(6):954–64. [https://](https://doi.org/10.1053/j.ajkd.2015.11.011)
 356 doi.org/10.1053/j.ajkd.2015.11.011

357

Summary of the case

1	PATIENT (GENDER, AGE)	Woman, 75 years old
2	FINAL DIAGNOSIS	Amoxicillin-induced crystal nephropathy
3	SYMPTOMS	General malaise, fever and painfully swollen lower left leg
4	MEDICATIONS	Amoxicillin high dose, after diagnosis changed to Penicillin G
5	CLINICAL PROCEDURE	Urine analysis revealing elongated crystals. Discontinuation of Amoxicillin and switch to Penicillin G.
6	SPECIALTY	Internal medicine